

DOUBLE CONSTRUCTION FOR CROSSED HOPF COALGEBRAS

MARCO ZUNINO

ABSTRACT. We provide an analog of the Drinfeld quantum double construction in the context of crossed Hopf group coalgebras introduced by Turaev. We prove that, provided the base group is finite, the double of a semisimple crossed Hopf group coalgebra is both modular and unimodular.

CONTENTS

Introduction	1
1. T-coalgebras	3
2. T-algebras	5
3. The outer dual and the inner dual of a T-coalgebra	6
4. Quasitriangular T-coalgebras	7
5. The quantum double of a finite-type T-coalgebra	8
6. Ribbon T-coalgebras	15
7. The quantum double of a semisimple T-coalgebra	16
8. The ribbon extension of a quasitriangular T-coalgebra	18
References	21

INTRODUCTION

Recently, Turaev [19, 20] (see also Le and Turaev [8] and Virelizier [23]) generalized the notion of a TQFT and Reshetikhin-Turaev invariants to the case of 3-manifolds endowed with a homotopy classes of maps to $K(\pi, 1)$, where π is a group. One of the key points in [19] is the notion of a *crossed Hopf π -coalgebra*, here called a *Turaev coalgebra* or, briefly, a *T-coalgebra* (see Section 1). As one can use categories of representations of modular Hopf algebras to construct Reshetikhin-Turaev invariants of 3-manifolds, one can use categories of representations of modular T-coalgebras to construct homotopy invariants of maps from 3-manifolds to $K(\pi, 1)$. Similarly, to construct Virelizier Hennings-like homotopy invariants [23], we need a ribbon T-coalgebra H such that the neutral component H_1 is unimodular [15].

Roughly speaking, a T-coalgebra H is a family $\{H_\alpha\}_{\alpha \in \pi}$ of algebras endowed with a *comultiplication* $\Delta_{\alpha, \beta}: H_{\alpha\beta} \rightarrow H_\alpha \otimes H_\beta$, a *counit* $\varepsilon: \mathbb{k} \rightarrow H_1$ (where 1 is the neutral element of π), and an *antipode* $s_\alpha: H_\alpha \rightarrow H_{\alpha^{-1}}$. It is also required that H is endowed with a family of algebra isomorphisms $\varphi_\beta^\alpha = \varphi_\beta: H_\alpha \rightarrow H_{\beta\alpha\beta^{-1}}$, the *conjugation*, compatible with the above structures and such that $\varphi_{\beta\gamma} = \varphi_\beta \circ \varphi_\gamma$. When $\pi = \{1\}$, we recover the definition of a Hopf algebra. A T-coalgebra H is of *finite type* when every H_α is finite-dimensional. H is *totally-finite* when the direct sum $\bigoplus_{\alpha \in \pi} H_\alpha$ is finite-dimensional. A *universal R-matrix* and a *twist*

Date: Friday, 13 September 2002.

2000 Mathematics Subject Classification. 81R50, 16W30, 57R56.

for a T-coalgebra H are, respectively, families $R = \{\xi_{(\alpha)} \otimes \zeta_{(\beta)} = R_{\alpha,\beta} \in H_\alpha \otimes H_\beta\}_{\alpha,\beta \in \pi}$ and $\theta = \{\theta_\alpha \in H_\alpha\}_{\alpha \in \pi}$ satisfying axioms that explicitly involve the conjugation (see, respectively, Sections 4 and 6). H is *modular* when it is ribbon and its component H_1 is modular (see Section 7). Properties of T-coalgebras are studied in [22].

Starting from a finite-dimensional Hopf algebra H , Drinfeld [2] showed how to obtain a quasitriangular Hopf algebra $D(H)$, the *quantum double of H* , such that the following conditions are satisfied (for details, see, e.g., [5]).

- There are embeddings of Hopf algebras $i: H \hookrightarrow D(H)$ and $j: H^{*\text{cop}} \hookrightarrow D(H)$.
- The linear map $H \otimes H^{*\text{cop}} \xrightarrow{i \otimes j} D(H) \otimes D(H) \xrightarrow{\mu} D(H)$ is bijective, where μ is the multiplication in $D(H)$.
- The universal R -matrix of $D(H)$ is the image of the canonical element of $H \otimes H^{*\text{cop}}$ under the embedding $i \otimes j: H \otimes H^{*\text{cop}} \hookrightarrow D(H) \otimes D(H)$.

In this paper we provide an analog of the quantum double for T-coalgebra H . We start (Section 5, Theorem 5.1) with an abstract definition of $D(H)$, as a solution of a universal problem, analogous to the above definition of the standard quantum double. The notion of a T-coalgebra is not self-dual, i.e., given a T-coalgebra $H = \{H_\alpha\}_{\alpha \in \pi}$, the family $H_\alpha^* = \{H_\alpha^*\}_{\alpha \in \pi}$ does not have a natural structure of a T-coalgebra. However, when H is of finite type, we introduce a T-coalgebra $H^{*\text{tot}}$, the *inner dual of H* , that, in many aspects, in particular in the construction of $D(H)$, plays the role of a dual for H (see Section 3). The components of $H^{*\text{tot}}$ are all isomorphic as algebras. In particular, as a vector space, $H_\alpha^{*\text{tot}} = \bigoplus_{\beta \in \pi} H_\beta^*$ for any $\alpha \in \pi$.

An explicit description of $D(H)$ is given in Theorems 5.3 and 5.6. In particular, every component $D_\alpha(H)$ of $D(H)$ is, as a vector space, $H_{\alpha^{-1}} \otimes \bigoplus_{\beta \in \pi} H_\beta^*$. So, in general, $D(H)$ is of finite type if and only if H is totally-finite and, in that case, $D(H)$ is also totally-finite. The product in $D_\alpha(H)$ is obtained by setting

$$(1_{\alpha^{-1}} \otimes f)(h \otimes \varepsilon) = h \otimes f \quad \text{and} \\ (h \otimes \varepsilon)(1_{\alpha^{-1}} \otimes f) = h''_{(\alpha^{-1})} \otimes \langle f, s_\gamma^{-1}(h'''_{(\gamma^{-1})}) \cdot \varphi_\alpha(h'_{(\alpha^{-1}\gamma\alpha)}) \rangle.$$

for any $f \in \bigoplus_{\beta \in \pi} H_\beta^*$, and $h \in H_{\alpha^{-1}}$. When $\pi = \{1\}$, we recover the standard definition of the quantum double of a Hopf algebra. We will see that both the product and the coproduct in $D(H)$ explicitly depend on the conjugation φ of H .

The quantum double $D(H)$ of a semisimple Hopf algebra H over a field of characteristic 0 is both semisimple [7, 11] and modular [3]. Here we prove that the double $D(H)$ of a semisimple T-coalgebra H over a field of characteristic 0 is semisimple if and only if H is totally-finite, and, in that case, $D(H)$ is also modular (Theorem 7.1). The key point in the proof is that, when H is totally finite, it gives rise to a graded Hopf algebra $H_{\text{pk}} = \bigoplus_{\alpha \in \pi} H_\alpha$, the *packed form of H* . The Hopf algebras $D(H_{\text{pk}})$ and $(D(H))_{\text{pk}}$ are different, but it is always possible to embed $D_1(H)$ in $D(H_{\text{pk}})$ as an algebra.

Last section is devoted to the generalization of the Reshetikhin-Turaev ribbon construction. Given a quasitriangular Hopf algebra H , Reshetikhin and Turaev [13] embedded it into a ribbon Hopf algebra $\text{RT}(H)$ that is a quotient of the polynomial algebra $H[\theta]$. Starting from a quasitriangular T-coalgebra H (not necessarily of finite type), we construct a ribbon T-coalgebra $\text{RT}(H)$ such that, when $\pi = \{1\}$, we recover the Reshetikhin-Turaev construction for Hopf algebras. As a corollary, for any T-coalgebra H of finite type (but not necessarily totally-finite), we obtain a ribbon T-coalgebra $\text{RT}(D(H))$.

The quantum double and the ribbon extension have categorical counterparts [4, 6, 16]. In [27], we provide a generalization of these constructions. In particular, the generalization of the center construction (corresponding to the quantum double in the algebraic case) looks very natural and as the simplest way to adapt the standard construction to the crossed case. Both the definition of $D(H)$ and the definition of $RT(H)$ introduced here are coherent to the categorical constructions as we briefly explain. In the standard case, starting from a tensor category $\mathcal{C}$, we obtain a braided tensor category $\mathcal{Z}(\mathcal{C})$ via the center construction [4]. Now, take $\mathcal{C} = \text{Rep}(H)$, the category of representations of a finite-dimensional Hopf algebra H . Both $\mathcal{Z}(\mathcal{C})$ and $\text{Rep}(D(H))$ are canonically isomorphic to the category of Yetter-Drinfeld modules over H (see [10, 24]). The corresponding result is still true when we consider a T-coalgebra H instead of an Hopf algebra, that is, also in the crossed case, we have a commutative diagram

$$\begin{array}{ccccc}
 \text{Rep}(H) & \xrightarrow{\mathcal{Z}} & \mathcal{Z}(\text{Rep}(H)) & \xlongequal{\quad} & \text{YD}(H) & \xlongequal{\quad} & \text{Rep}(D(H)) \\
 \uparrow \text{Rep} & & & & & & \uparrow \text{Rep} \\
 H & \xrightarrow{\quad D \quad} & & & & & D(H)
 \end{array}$$

(where, of course, we also need to generalize the notions of a tensor category and of a Yetter-Drinfeld module). Similar results are obtained for the ribbon construction.

ACKNOWLEDGMENTS. The author wants to thank his adviser V. Turaev for his stimulating and constructive direction in the research. The author also wants to thank A. Bruguières, B. Enriquez, Ch. Kassel, H.J. Schneider, L. Vainerman, and A. Virelizier for their useful remarks.

The author was partially supported by the INDAM, ISTITUTO NAZIONALE DI ALTA MATEMATICA, Rome.

1. T-COALGEBRAS

We recall the notion of a T-coalgebra as in [20]. T-coalgebras are a generalization of standard notion Hopf algebras (for a general reference to Hopf algebras, see [17, 1] or, for a modern introduction, [15]). A generalization of the Heynemann-Sweedler notation [17] is also provided. At the end of this section, we study some properties of the antipode of a T-coalgebra.

BASIC DEFINITIONS. Let $\mathbb{k}$ be a commutative field and let π be a group. A T-coalgebra H (over π and $\mathbb{k}$) is given by the following data.

- For any $\alpha \in \pi$, an associative $\mathbb{k}$ -algebra H_α , called the α -th component of H . The multiplication is denoted $\mu_\alpha: H_\alpha \otimes H_\alpha \rightarrow H_\alpha$ and the unit is denoted $\eta_\alpha: \mathbb{k} \rightarrow H_\alpha$, with $1_\alpha = \eta_\alpha(1)$.
- A family of algebra morphisms $\Delta = \{\Delta_{\alpha,\beta}: H_{\alpha\beta} \rightarrow H_\alpha \otimes H_\beta\}_{\alpha,\beta \in \pi}$, called *comultiplication*, that is *coassociative* in the sense that, for any $\alpha, \beta, \gamma \in \pi$, we have $(H_\alpha \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma} = (\Delta_{\alpha,\beta} \otimes H_\gamma) \circ \Delta_{\alpha\beta,\gamma}$.
- An algebra morphism $\varepsilon: H_1 \rightarrow \mathbb{k}$, called *counit*, such that, for any $\alpha \in \pi$, we have $(\varepsilon \otimes H_\alpha) \circ \Delta_{1,\alpha} = \text{Id}$ and $(H_\alpha \otimes \varepsilon) \circ \Delta_{\alpha,1} = \text{Id}$.
- A set of algebra isomorphisms $\varphi = \{\varphi_\beta^\alpha: H_\alpha \rightarrow H_{\beta\alpha\beta^{-1}}\}_{\alpha,\beta \in \pi}$ called *conjugation*. When not strictly necessary, the upper index will be omitted. We require that φ satisfies the following conditions.
 - φ is *multiplicative*, i.e., $\varphi_\beta \circ \varphi_\gamma = \varphi_{\beta\gamma}$, for any $\beta, \gamma \in \pi$. It follows that $\varphi_1^\alpha = \text{Id}$, for any $\alpha \in \pi$.
 - φ is *compatible with Δ* , i.e., $\Delta_{\gamma\alpha\gamma^{-1}, \gamma\beta\gamma^{-1}} \circ \varphi_\gamma = (\varphi_\gamma \otimes \varphi_\gamma) \circ \Delta_{\alpha,\beta}$, for any $\alpha, \beta, \gamma \in \pi$.

- φ is *compatible with ε* , i.e., $\varepsilon \circ \varphi_\gamma = \varepsilon$ for any $\gamma \in \pi$.
- Finally, a set of $\mathbb{k}$ -linear morphisms $s = \{s_\alpha: H_\alpha \rightarrow H_{\alpha^{-1}}\}_{\alpha \in \pi}$, the *antipode*, such that, for any $\alpha \in \pi$,

$$(1) \quad \mu_\alpha \circ (s_{\alpha^{-1}} \otimes H_\alpha) \circ \Delta_{\alpha^{-1}, \alpha} = \eta_\alpha \circ \varepsilon = \mu_\alpha \circ (H_\alpha \otimes s_{\alpha^{-1}}) \circ \Delta_{\alpha, \alpha^{-1}}.$$

(The compatibility of the antipode with the conjugation isomorphisms is a consequence of the axioms.) In [20], a T-coalgebra is called a *crossed group Hopf coalgebra*.

We say that H is of *finite-type* when any component H_α (with $\alpha \in \pi$) is a finite-dimensional $\mathbb{k}$ -vector space. We say that H is *totally-finite* when $\dim_{\mathbb{k}} \bigoplus_{\alpha \in \pi} H_\alpha < \infty$, i.e., when H is of finite-type and almost all the H_α are zero. It was proved in [22] that the antipode of a finite-type T-coalgebra is always bijective.

We observe that the component H_1 of a T-coalgebra H is a Hopf algebra in the usual sense. We also observe that, for $\pi = \{1\}$, we recover the usual notion of a Hopf algebra.

Example 1.1 (TH-coalgebras). Let H_1 be a Hopf algebra, with comultiplication Δ_1 , counit ε_1 , and antipode s_1 , endowed with a group morphism

$$\varphi^1: \pi \rightarrow \text{Aut}(H_1): \alpha \mapsto \varphi_\alpha^1,$$

where $\text{Aut}(H_1)$ is the group of Hopf algebra automorphisms of H_1 . We obtain a T-coalgebra H by setting (for any $\alpha, \beta \in \pi$) $H_\alpha = H_1$ as algebra, $\Delta_{\alpha, \beta} = \Delta_1$, $\varepsilon = \varepsilon_1$, $s_\alpha = s_1$, and $\varphi_\beta^\alpha = \varphi_\beta^1: H_\alpha \rightarrow H_{\beta\alpha\beta^{-1}}$. The T-coalgebra H is called the *TH-coalgebra based on H_1* (the H stand for Hopf, since H is also an Hopf algebra).

COOPPOSITE T-COALGEBRA. Let H be a T-coalgebra with invertible antipode. The *coopposite T-coalgebra* H^{cop} is the T-coalgebra defined as follows.

- For any $\alpha \in \pi$, we set $H_\alpha^{\text{cop}} = H_{\alpha^{-1}}$ as an algebra.
- The comultiplication Δ^{cop} is obtained by setting, for any $\alpha, \beta \in \pi$,

$$\Delta_{\alpha, \beta}^{\text{cop}} = \left(H_{\alpha\beta}^{\text{cop}} = H_{\beta^{-1}\alpha^{-1}} \xrightarrow{\Delta_{\beta^{-1}, \alpha^{-1}}} H_{\beta^{-1}} \otimes H_{\alpha^{-1}} \xrightarrow{\sigma} H_{\alpha^{-1}} \otimes H_{\beta^{-1}} = H_\alpha^{\text{cop}} \otimes H_\beta^{\text{cop}} \right).$$

Δ^{cop} has a counit $\varepsilon^{\text{cop}} = \varepsilon$.

- The antipode s^{cop} is obtained by setting $s_\alpha^{\text{cop}} = s_\alpha^{-1}$, for any $\alpha \in \pi$.
- The conjugation φ^{cop} is obtained by setting $\varphi_\beta^{\text{cop}} = \varphi_\beta$, for any $\beta \in \pi$.

In particular, when H is a TH-coalgebra, H^{cop} is the TH-coalgebra based on H_1^{cop} . **HEYNEMANN-SWEEDLER NOTATION.** The coassociativity of H allows us to introduce an analog of the Heynemann-Sweedler notation, see [17]. Given $\alpha_1, \dots, \alpha_n \in \pi$ and defining

$$\Delta_{\alpha_1, \alpha_2, \dots, \alpha_n} = \left(H_{\alpha_1 \alpha_2 \dots \alpha_n} \xrightarrow{\Delta_{\alpha_1, \alpha_2 \dots \alpha_n}} H_{\alpha_1} \otimes H_{\alpha_2 \dots \alpha_n} \xrightarrow{H_1 \otimes \Delta_{\alpha_2, \alpha_3 \dots \alpha_n}} \dots \rightarrow H_{\alpha_1} \otimes H_{\alpha_2} \otimes \dots \otimes H_{\alpha_n} \right),$$

for any $h \in H_{\alpha_1 \alpha_2 \dots \alpha_n}$ we set

$$h'_{(\alpha_1)} \otimes h''_{(\alpha_2)} \otimes \dots \otimes h'^n_{(\alpha_n)} = \Delta_{\alpha_1, \alpha_2, \dots, \alpha_n}(h).$$

Let M be a vector space over $\mathbb{k}$ and suppose that $f: H_{\alpha_1} \times H_{\alpha_2} \times \dots \times H_{\alpha_n} \rightarrow M$ is a $\mathbb{k}$ -multilinear map. Let $\hat{f}$ denote the linear map $H_{\alpha_1} \otimes H_{\alpha_2} \otimes \dots \otimes H_{\alpha_n} \rightarrow M$ induced by f . We introduce the notation

$$f(h'_{(\alpha_1)}, h''_{(\alpha_2)}, \dots, h'^n_{(\alpha_n)}) = \hat{f}(\Delta_{\alpha_1, \alpha_2, \dots, \alpha_n}(h)).$$

For simplicity, we also suppress the subscript (α_i) when $\alpha_i = 1$.

PROPERTIES OF THE ANTIPODE. Let H be a T-coalgebra and let A be an algebra with multiplication μ_A and unit η_A . We define a *convolution algebra* $\text{Conv}(H, A)$ (see [22]) in the following way. As a vector space, $\text{Conv}(H, A) = \bigoplus_{\beta \in \pi} \text{Hom}_{\mathbb{k}}(H_{\beta}, A)$. The multiplication in $\text{Conv}(H, A)$ is obtained by setting, for any $\beta_1, \beta_2 \in \pi$, $f_1 \in \text{Hom}_{\mathbb{k}}(H_{\beta_1}, A)$, $f_2 \in \text{Hom}_{\mathbb{k}}(H_{\beta_2}, A)$, and $h \in H_{\beta_1 \beta_2}$,

$$(f_1 * f_2)(h) = f_1(h'_{(\beta_1)}) f_2(h''_{(\beta_2)}).$$

With this multiplication, $\text{Conv}(H, A)$ becomes an associative algebra.

For any $\alpha \in \pi$, we introduce the notation $\text{Conv}_{\alpha}(H) = \text{Conv}(H, H_{\alpha})$. It is clear that (1) is equivalent to say that $s_{\alpha^{-1}}$ is a two-sided inverse of the identity morphism of H_{α} in the convolution algebra $\text{Conv}_{\alpha}(H)$. Thus, we can reformulate the axiom for the antipode of H by requiring that, for any $\alpha \in \pi$, the identity morphism of H_{α} is invertible in $\text{Conv}_{\alpha}(H)$. This also proves that the antipode of a T-coalgebra is unique.

Lemma 1.2. *Let H be a T-coalgebra. The antipode s of H is both antimultiplicative and anticomultiplicative, i.e.,*

$$(2) \quad s_{\alpha}(hk) = s_{\alpha}(k)s_{\alpha}(h)$$

for any $\alpha \in \pi$, $h, k \in H_{\alpha}$, and

$$(3) \quad s_{\alpha}(h'_{(\alpha)}) \otimes s_{\alpha}(h''_{(\beta)}) = (s_{\alpha\beta}(h))''_{(\alpha^{-1})} \otimes (s_{\alpha\beta}(h))'_{(\beta^{-1})}$$

for any $h \in H_{\alpha\beta}$.

The proof can be obtained as in the standard case (i.e., for $\pi = \{1\}$).

2. T-ALGEBRAS

When we dualize the axioms of a T-coalgebra, we obtain the notion of a T-algebra. A T-algebra H can be equivalently described as a Hopf algebra endowed with a family of automorphisms, the *packed form of H* . In this section, we present both approaches.

BASIC DEFINITIONS. A T-algebra H is a family $\{(H_{\alpha}, \Delta_{\alpha}, \eta_{\alpha})\}_{\alpha \in \pi}$ of $\mathbb{k}$ -coalgebras, endowed with the following data.

- A family of coalgebra morphisms $\mu = \{\mu_{\alpha, \beta}: H_{\alpha} \otimes H_{\beta} \rightarrow H_{\alpha\beta}\}_{\alpha, \beta \in \pi}$, called *multiplication*, that is *associative*, in the sense that, for any $\alpha, \beta, \gamma \in \pi$,

$$(4) \quad \mu_{\alpha\beta, \gamma} \circ (\mu_{\alpha, \beta} \otimes H_{\gamma}) = \mu_{\alpha, \beta\gamma} \circ (H_{\alpha} \otimes \mu_{\beta, \gamma}).$$

Given $h \in H_{\alpha}$ and $k \in H_{\beta}$, with $\alpha, \beta \in \pi$, we set $hk = \mu_{\alpha, \beta}(h, k)$. With this notation, (4) can be simply rewritten as $(hk)l = h(kl)$ for any $h \in H_{\alpha}$, $k \in H_{\beta}$, $l \in H_{\gamma}$ and $\alpha, \beta, \gamma \in \pi$.

- An algebra morphism $\eta: \mathbb{k} \rightarrow H_1$, called *unit*, such that, if we set $1 = \eta(1_{\mathbb{k}})$, then, for any $h \in H_{\alpha}$ (with $\alpha \in \pi$), we have $1h = h = h1$.
- A set of coalgebra isomorphism $\psi = \{\psi_{\beta}^{\alpha}: H_{\alpha} \rightarrow H_{\beta\alpha\beta^{-1}}\}_{\alpha, \beta \in \pi}$, called *conjugation*. Also in this case, when not strictly necessary, the upper index will be omitted. We require that ψ satisfies the following conditions.
 - ψ is *multiplicative*, i.e., for any α, β , and $\gamma \in \pi$, we have $\psi_{\beta} \circ \psi_{\gamma} = \psi_{\beta\gamma}$. It follows that, for any $\alpha \in \pi$, we have $\psi_1^{\alpha} = \text{Id}$.
 - ψ is *compatible with μ* , i.e., for any $\beta \in \pi$, we have $\psi_{\beta}(hk) = \psi_{\beta}(h)\psi_{\beta}(k)$.
 - ψ is *compatible with η* , i.e., for any $\beta \in \pi$, we have $\psi_{\beta}(1) = 1$.

- Finally, a set of linear isomorphisms $S = \{S_{\alpha}: H_{\alpha} \rightarrow H_{\alpha^{-1}}\}_{\alpha \in \pi}$, the *antipode*, such that, for any $\alpha \in \pi$, we have $\mu_{\alpha^{-1}, \alpha} \circ (S_{\alpha} \otimes H_{\alpha}) \circ \Delta_{\alpha} = \eta \circ \varepsilon_{\alpha} = \mu_{\alpha, \alpha^{-1}} \circ (H_{\alpha} \otimes S_{\alpha}) \circ \Delta_{\alpha}$.

For any $\alpha \in \pi$, the coalgebra H_{α} is called the α -th component of H .

PACKED FORM OF A T-ALGEBRA. Let H be a T-algebra. We define a Hopf algebra H_{pk} , that we call the *packed form of H* , as follows. As a coalgebra, H_{pk} is the direct sum of the components of H . For any $\alpha \in \pi$, we denote by i_α the inclusion of H_α in H_{pk} . The multiplication μ_{pk} of H_{pk} is the colimit, in the category of vector spaces, $\varinjlim_{\alpha, \beta \in \pi} (i_{\alpha\beta} \circ \mu_{\alpha, \beta})$, i.e., the only $\mathbb{k}$ -linear map from $H_{\text{pk}} \otimes H_{\text{pk}}$ to H_{pk} such that the restriction on $H_\alpha \otimes H_\beta \subset H_{\text{pk}} \otimes H_{\text{pk}}$ coincides with $\mu_{\alpha, \beta}$,

$$\begin{array}{ccc} H_\alpha \otimes H_\beta & \xrightarrow{\quad} & \bigoplus_{\gamma, \delta \in \pi} (H_\gamma \otimes H_\delta) = \left(\bigoplus_{\gamma \in \pi} H_\gamma \right) \otimes \left(\bigoplus_{\delta \in \pi} H_\delta \right) = H_{\text{pk}} \otimes H_{\text{pk}} \\ \mu_{\alpha, \beta} \downarrow & & \downarrow \mu_{\text{pk}} \\ H_{\alpha\beta} & \xrightarrow{\quad} & H_{\text{pk}} \end{array}.$$

The unit η_{pk} of H_{pk} is given by $i_1 \circ \eta_H$, i.e., $1_{\text{pk}} = 1 \in H_1 \subset H_{\text{pk}}$. Finally, the antipode S_{pk} of H_{pk} is given by the sum $\sum_{\alpha \in \pi} S_\alpha$. The Hopf algebra H_{pk} is endowed with a group morphism $\psi: \pi \rightarrow \text{Aut}(H_{\text{pk}}): \alpha \mapsto \psi_{\text{pk}, \alpha}$, where $\psi_{\text{pk}, \alpha} = \sum_{\beta \in \pi} \psi_\alpha^\beta: \bigoplus_{\beta \in \pi} H_\beta \mapsto \bigoplus_{\beta \in \pi} H_\beta$.

Conversely, let H_{tot} be a Hopf algebra with multiplication μ_{tot} , unit 1, and antipode S_{tot} , endowed with a group homomorphism $\varphi_{\text{tot}}: \pi \rightarrow \text{Aut}(H_{\text{tot}}): \alpha \mapsto \varphi_{\text{tot}, \alpha}$. Suppose that the following conditions are satisfied.

- There exists a family of sub-coalgebras $\{H_\alpha\}_{\alpha \in \pi}$ of H_{tot} such that $H_{\text{tot}} = \bigoplus_{\alpha \in \pi} H_\alpha$.
- $H_\alpha \cdot H_\beta \subset H_{\alpha\beta}$ for any $\alpha, \beta \in \pi$.
- $1 \in H_1$.
- For any $\alpha, \beta \in \pi$, $\varphi_{\text{tot}, \beta}$ sends $H_\alpha \subset H_{\text{tot}}$ to $H_{\beta\alpha\beta^{-1}} \subset H_{\text{tot}}$.
- For any $\alpha \in \pi$, we have $S_{\text{tot}}(H_\alpha) = H_{\alpha^{-1}}$.

Then, we obtain, in the obvious way, a T-algebra H such that $H_{\text{pk}} = H_{\text{tot}}$.

3. THE OUTER DUAL AND THE INNER DUAL OF A T-COALGEBRA

We study how to provide a convenient notion of a dual for a finite-type T-coalgebra H . The easiest way is to define a T-algebra H^* , the *outer dual of H* . However, for many purposes, in particular in the construction of the quantum double of H , it is convenient to introduce a TH-coalgebra $H^{*\text{tot}}$ based on the packed form H_{pk}^* of H^* , the *inner dual of H* .

THE OUTER DUAL. Let H be a finite-type T-coalgebra. The *outer dual of H* is the T-algebra H^* defined as follows. For any $\alpha \in \pi$, the α -th component of H^* is the dual coalgebra H_α^* of the algebra H_α . The multiplication of H^* is given by

$$(5) \quad \langle \mu_{\alpha, \beta}(f, g), h \rangle = \langle f \otimes g, \Delta_{\alpha, \beta}(h) \rangle$$

for any $f \in H_\alpha^*$, $g \in H_\beta^*$ and $h \in H_{\alpha\beta}$, with $\alpha, \beta \in \pi$. The unit of H^* is given by $\varepsilon \in H_1^* \subset H^*$. The antipode S^* of H^* is given by $S_\alpha^* = s_{\alpha^{-1}}^*$, for any $\alpha \in \pi$. Finally, for any $\beta \in \pi$, the conjugation isomorphism ψ_β^* of H^* is given by $\psi_\beta^* = \varphi_{\beta^{-1}}^*$.

THE INNER DUAL. Since $(H^*)_{\text{pk}}$ is a Hopf algebra endowed with a group homomorphism $\psi_{(H^*)_{\text{pk}}}: \pi \rightarrow \text{Aut}((H^*)_{\text{pk}})$, we can construct the TH-coalgebra based on $(H^*)_{\text{pk}}$. We call this TH-coalgebra the *inner dual of H* and we denote it $H^{*\text{tot}}$. Explicitly, $H_1^{*\text{tot}} = (H^*)_{\text{pk}}$ is obtained as follows.

- As a coalgebra, $H_1^{*\text{tot}} = \bigoplus_{\alpha \in \pi} H_\alpha^*$.
- The multiplication is obtained by (5), extending by linearity. The unit is given by $\varepsilon^{*\text{tot}} = \varepsilon \in H_1^* \subset \bigoplus_{\alpha \in \pi} H_\alpha^*$.
- The antipode is given by $s_1^{*\text{tot}} = \sum_{\alpha \in \pi} S_\alpha^* = \sum_{\alpha \in \pi} s_{\alpha^{-1}}^*$.
- Finally, $\psi_{H_{\text{pk}}^*, \beta} = \sum_{\beta \in \pi} \varphi_{\beta^{-1}}^*$.

THE COOPPOSITE INNER DUAL. Given any T-coalgebra H , then $((H^*)_{\text{pk}})^{\text{cop}}$ is the Hopf algebra obtained from $(H^*)_{\text{pk}}$ by replacing its comultiplication with the new one $\Delta_* = \Delta^{*\text{tot}, \text{cop}}$ given by

$$(6) \quad \langle \Delta_*(f), h \otimes k \rangle = \langle f, kh \rangle$$

for any $f \in H_\alpha^* \subset \bigoplus_{\beta \in \pi} H_\beta^*$ and $h, k \in H_\alpha$, with $\alpha \in \pi$. We also need to replace the antipode with the new one given by $s_* = S^{*, \text{cop}} = (S^*)^{-1}$. In particular, we have $\langle s_*(f), h \rangle = \langle f, s_\alpha^{-1}(h) \rangle$, for any $f \in H_\alpha^*$ and $h \in H_{\alpha^{-1}}$, with $\alpha \in \pi$. The TH-coalgebra based on $((H^*)_{\text{pk}})^{\text{cop}}$ is called the *coopposite inner dual of H* and is denoted $H^{*\text{tot}, \text{cop}}$. Notice that $\varphi_{H^{*\text{tot}, \text{cop}}, \alpha} = \varphi_{H^*, \alpha} = \sum_{\beta \in \pi} \varphi_{\beta^{-1}}^*$, for any $\alpha \in \pi$.

4. QUASITRIANGULAR T-COALGEBRAS

The notion of a quasitriangular Hopf algebra [2] is generalized to the case of a T-coalgebra in [20]. A *quasitriangular T-coalgebras* is a T-coalgebra H endowed with a family $R = \{R_{\alpha, \beta} = \xi_{(\alpha).i} \otimes \zeta_{(\beta).i} \in H_\alpha \otimes H_\beta\}_{\alpha, \beta \in \pi}$, called *universal R -matrix*, such that $R_{\alpha, \beta}$ is invertible for any $\alpha, \beta \in \pi$ and the following conditions are satisfied.

- For any $\alpha, \beta \in \pi$ and $h \in H_{\alpha\beta}$,

$$(7a) \quad R_{\alpha, \beta} \Delta_{\alpha, \beta}(h) = (\sigma \circ (\varphi_{\alpha^{-1}} \otimes H_\alpha) \circ \Delta_{\alpha\beta\alpha^{-1}, \alpha})(h) R_{\alpha, \beta}.$$

- For any $\alpha, \beta, \gamma \in \pi$,

$$(7b) \quad (H_\alpha \otimes \Delta_{\beta, \gamma})(R_{\alpha, \beta\gamma}) = (R_{\alpha, \gamma})_{1\beta 3} (R_{\alpha, \beta})_{12\gamma},$$

where, given two vector spaces P and Q over $\mathbb{k}$, for any $x = p_i \otimes q_i \in P \otimes Q$ we set $x_{1\beta 3} = p_i \otimes 1_\beta \otimes q_i$, and $x_{12\gamma} = p_i \otimes q_i \otimes 1_\gamma$.

- For any $\alpha, \beta, \gamma \in \pi$,

$$(7c) \quad (\Delta_{\alpha, \beta} \otimes H_\gamma)(R_{\alpha\beta, \gamma}) = ((\varphi_\beta \otimes H_\gamma)(R_{\beta^{-1}\alpha\beta, \gamma}))_{1\beta 3} (R_{\beta, \gamma})_{\alpha 23},$$

where, given two vector spaces P and Q , for any $x = p_i \otimes q_i \in P \otimes Q$ we set $x_{\alpha 23} = 1_\alpha \otimes p_i \otimes q_i$.

- R is *compatible with φ* , in the sense that, for any $\alpha, \beta, \gamma \in \pi$, we have

$$(7d) \quad (\varphi_\alpha \otimes \varphi_\alpha)(R_{\beta, \gamma}) = R_{\alpha\beta\alpha^{-1}, \alpha\gamma\alpha^{-1}}.$$

Notice that $(H_1, R_{1,1})$ is a quasitriangular Hopf algebra in the usual sense.

For any $\alpha, \beta \in \pi$, we introduce the notation $\tilde{\xi}_{(\alpha).i} \otimes \tilde{\zeta}_{(\beta).i} = \tilde{R}_{\alpha, \beta} = (R^{-1})_{\alpha, \beta}$.

Remark 4.1 (Yang-Baxter equation). It is proved in [20] that, for any $\alpha, \beta, \gamma \in \pi$ we have

$$(8) \quad (R_{\beta, \gamma})_{\alpha 23} (R_{\alpha, \gamma})_{1\beta 3} (R_{\alpha, \beta})_{12\gamma} \\ = (R_{\alpha, \beta})_{12\gamma} ((H_\alpha \otimes \varphi_{\beta^{-1}})(R_{\alpha, \beta\gamma\beta^{-1}}))_{1\beta 3} (R_{\beta, \gamma})_{\alpha 23}.$$

This is an analog of the standard Yang-Baxter equation (see, e.g., [5]).

THE MIRROR T-COALGEBRA. Let $H = (H, R)$ be a quasitriangular Hopf algebra (with $R = \xi_i \otimes \zeta_i$ and $R^{-1} = \tilde{R} = \tilde{\xi}_i \otimes \tilde{\zeta}_i$). By replacing R with $\bar{R} = \sigma(\tilde{R}) = \tilde{\zeta}_i \otimes \tilde{\xi}_i$, we obtain another quasitriangular structure $\bar{H} = (H, \bar{R})$. This means that, in the category of representations of H , we replace the braiding c_R provided by R by the braiding $c_{\bar{R}}^{-1}$ provided by $\bar{R}$. When H is a T-coalgebra, the family $\{R_{\alpha, \beta}^{-1} = \tilde{\zeta}_{\alpha.i} \otimes \tilde{\xi}_{\beta.i}\}_{\alpha, \beta \in \pi}$ is not a universal R -matrix for H . Nevertheless, it is still possible to generalize the definition of $\bar{H}$ in the following way [20].

Let H be a T-coalgebra. The T-coalgebra $\bar{H}$, called *the mirror of H* [20], is defined as follows.

- For any $\alpha \in \pi$, we set $\bar{H}_\alpha = H_{\alpha^{-1}}$.

- For any $\alpha, \beta \in \pi$, the component $\overline{\Delta}_{\alpha, \beta}$ of the comultiplication $\overline{\Delta}$ of $\overline{H}$ is given by

$$(9) \quad \overline{\Delta}_{\alpha, \beta}(h) = ((\varphi_\beta \otimes H_{\beta^{-1}}) \circ \Delta_{\beta^{-1}\alpha\beta, \beta^{-1}})(h) \in H_{\alpha^{-1}} \otimes H_{\beta^{-1}} = \overline{H}_\alpha \otimes \overline{H}_\beta,$$

for any $h \in H_{\beta^{-1}\alpha^{-1}} = \overline{H}_{\alpha\beta}$. If we set $h'_{(\alpha)} \otimes h''_{(\beta)} = \overline{\Delta}_{\alpha, \beta}(h)$, then (9) can be written in the form $h'_{(\alpha)} \otimes h''_{(\beta)} = \varphi_\beta(h'_{(\beta^{-1}\alpha^{-1}\beta)}) \otimes h''_{(\beta^{-1})}$. The counit of $\overline{H}$ is given by $\varepsilon \in H_1^* = \overline{H}_1^*$.

- For any $\alpha \in \pi$, the α -th component of the antipode $\overline{s}$ of $\overline{H}$ is given by $\overline{s}_\alpha = \varphi_\alpha \circ s_{\alpha^{-1}}$.
- Finally, for any $\alpha \in \pi$, we set $\overline{\varphi}_\alpha = \varphi_\alpha$.

If H is quasitriangular, then $\overline{H}$ is also quasitriangular with universal R -matrix $\overline{R}$ given by

$$(10) \quad \overline{R}_{\alpha, \beta} = (\sigma(R_{\beta^{-1}, \alpha^{-1}}))^{-1} \in H_{\alpha^{-1}} \otimes H_{\beta^{-1}} = \overline{H}_\alpha \otimes \overline{H}_\beta$$

for any $\alpha, \beta \in \pi$.

If, for any $\alpha, \beta \in \pi$, we introduce the notation $\overline{\xi}_{(\alpha).i} \otimes \overline{\zeta}_{(\beta).i} = \overline{R}_{\alpha, \beta}$, then we can write (10) in the form $\overline{\xi}_{(\alpha).i} \otimes \overline{\zeta}_{(\beta).i} = \tilde{\zeta}_{(\alpha^{-1}).i} \otimes \tilde{\xi}_{(\beta^{-1}).i}$.

Notice that $\overline{\overline{H}} = H$. Notice also that, due to the definition of $\overline{\Delta}$, the mirror of a TH-coalgebra is not, in general, a TH-coalgebra. In particular, the mirror of the inner dual of a finite-type T-coalgebra is not a TH-coalgebra.

5. THE QUANTUM DOUBLE OF A FINITE-TYPE T-COALGEBRA

Theorem 5.1. *Let H be a finite-type T-coalgebra. There exists a unique quasitriangular T-coalgebra $D(H)$, the quantum double of H , such that the following conditions are satisfied.*

- The T-coalgebras $\overline{H}$ and $H^{*\text{tot}, \text{cop}}$ can be embedded into $D(H)$, i.e., there are two morphisms of T-coalgebras $i: \overline{H} \rightarrow D(H)$ and $j: H^{*\text{tot}, \text{cop}} \rightarrow D(H)$ such that i_α and j_α are injective for any $\alpha \in \pi$.
- For any $\alpha \in \pi$, the linear map

$$(11) \quad p_\alpha = \left(H_1^{*\text{tot}, \text{cop}} \otimes \overline{H}_\alpha \xrightarrow{j_\alpha \otimes i_\alpha} D_\alpha(H) \otimes D_\alpha(H) \xrightarrow{\mu_\alpha} D_\alpha(H) \right)$$

is bijective (where $D_\alpha(H)$ is the α -th component of $D(H)$ and μ_α is the multiplication in $D_\alpha(H)$).

- For any $\alpha, \beta \in \pi$, the component $R_{\alpha, \beta}$ of the R -matrix R of $D(H)$ is the image of the canonical element of $H_{\alpha^{-1}} \otimes H_1^{*\text{tot}, \text{cop}}$ under the embedding $j_\alpha \otimes i_\beta: H_1^{*\text{tot}, \text{cop}} \otimes H_{\alpha^{-1}} \hookrightarrow D_\alpha(H) \otimes D_\beta(H)$.

To prove Theorem 5.1, firstly construct a quasitriangular T-coalgebra $D(H)$. Then, we will see that $D(H)$ is a solution of the above universal problem.

Remark 5.2. Notice that, when $\pi = \{1\}$, both $D(H)$ and $D(\overline{H})$ coincide with the standard definition of the quantum double of H . The reason of the convention that takes the mirror of H in the above description is that we want to be coherent with the conventions in [6]. However, often in the standard case p_1 is defined as $\mu_1 \circ (i_1 \otimes j_1)$, reversing the position of i_1 and j_1 . Moreover, some authors identify $D_1(H)$ with the vector space $H_1^* \otimes H$. However, since on that point it seems there is no standard convention, here we also follow the notations in [6], apart for the detail that we reversed the order of the factors in the tensor product.

CONSTRUCTION OF $D(H)$. Let H be a finite-type T-coalgebra. We realize $D(H)$ as follows.

- For any $\alpha \in \pi$, the α -th component of $D(H)$, denoted $D_\alpha(H)$, is, as a vector space,

$$H_{\alpha^{-1}} \otimes H_\alpha^{*\text{tot},\text{cop}} = H_{\alpha^{-1}} \otimes H_1^{*\text{tot},\text{cop}} = H_\alpha \otimes \bigoplus_{\beta \in \pi} H_\beta^*.$$

The multiplication in $D_\alpha(H)$ is not obtained by tensor product of algebras of $H_{\alpha^{-1}}$ and $H_\alpha^{*\text{tot},\text{cop}}$. In the sequel, given $h \in H_{\alpha^{-1}}$ and $F \in H_\alpha^{*\text{tot},\text{cop}}$, the element in $D(H)$ corresponding to $h \otimes F$ in $D(H)$ will be denoted $h \otimes F$.

In view of the role played by $H^{*\text{tot},\text{cop}}$ in the construction of the quantum double, the Heynemann-Sweedler notation will be reserved for the comultiplication of $H^{*\text{tot},\text{cop}}$, not of $H^{*\text{tot}}$, i.e., given $F \in \sum_{\alpha \in \pi} H_\alpha^*$, we set $F' \otimes F'' = \Delta_*(F)$.

Let f be in H_α^* and let h and k be in H_α , with $\alpha \in \pi$. By $\langle f, h \cdot k \rangle$ we denote the linear functional on H_α that evaluated at $x \in H_\alpha$ gives $\langle f, h x k \rangle$. $D_\alpha(H)$ is an algebra under the multiplication obtained by setting, for any $h, k \in H_{\alpha^{-1}}$, $f \in H_\gamma^*$, and $g \in H_\delta^*$, with $\gamma, \delta \in \pi$,

$$(12) \quad (h \otimes f)(k \otimes g) = h''_{(\alpha^{-1})} k \otimes f \langle g, s_\delta^{-1}(h'''_{(\delta^{-1})}) \cdot \varphi_\alpha(h'_{(\alpha^{-1}\delta\alpha)}) \rangle.$$

The unit of $D_\alpha(H)$ is given by $1_{\alpha^{-1}} \otimes \varepsilon$. It follows that the canonical embeddings $H_{\alpha^{-1}}, H_\alpha^{*\text{tot},\text{cop}} \hookrightarrow D_\alpha(H)$ are algebra morphisms and that, for any $h \in H_{\alpha^{-1}}$ and $f \in H_\gamma^*$, we have

$$(13a) \quad (1_{\alpha^{-1}} \otimes f)(h \otimes \varepsilon) = h \otimes f \quad \text{and}$$

$$(13b) \quad (h \otimes \varepsilon)(1_{\alpha^{-1}} \otimes f) = h''_{(\alpha^{-1})} \otimes \langle f, s_\gamma^{-1}(h'''_{(\gamma^{-1})}) \cdot \varphi_\alpha(h'_{(\alpha^{-1}\gamma\alpha)}) \rangle.$$

- The comultiplication is given by

$$(14) \quad \Delta_{\alpha,\beta}(h \otimes F) = \left(\varphi_\beta(h'_{(\beta^{-1}\alpha^{-1}\beta)}) \otimes F' \right) \otimes \left(h''_{(\beta^{-1})} \otimes F'' \right),$$

for any $\alpha, \beta \in \pi$, $h \in \overline{H}_{\alpha\beta} = H_{\beta^{-1}\alpha^{-1}}$, and $F \in H_{\alpha\beta}^{*\text{tot},\text{cop}}$ (We recall that $F' \otimes F'' = \Delta_*(F)$, see (6)). The counit is obtained by setting $\langle \varepsilon, h \otimes f \rangle = \langle \varepsilon, h \rangle \langle f, 1_\gamma \rangle$, for any $h \in H_1$ and $f \in H_\gamma^*$, with $\gamma \in \pi$.

- For any $\alpha \in \pi$, the α -th component of the antipode of $D(H)$ is given by

$$(15) \quad s_\alpha(h \otimes F) = (\overline{s}_\alpha(h) \otimes \varepsilon)(1 \otimes s_*(F)) = ((\varphi_\alpha \circ s_{\alpha^{-1}})(h) \otimes \varepsilon)(1_\alpha \otimes s_*(F)),$$

for any $h \in \overline{H}_\alpha = H_{\alpha^{-1}}$ and $F \in H_\alpha^{*\text{tot},\text{cop}}$, where s_* is the antipode of $H^{*\text{tot},\text{cop}}$ and $\overline{s}_\alpha = \varphi_\alpha \circ s_{\alpha^{-1}}$ is the antipode of $\overline{H}$.

- Finally, for any $\alpha \in \pi$, we set

$$(16) \quad \varphi_\beta(h \otimes f) = \varphi_\beta(h) \otimes \varphi_{H^{*\text{tot},\text{cop}},\beta}(f) = \varphi_\beta(h) \otimes \varphi_{\beta^{-1}}^*(f),$$

for any $h \in H_{\alpha^{-1}}$ and $f \in H_\gamma^{*\text{tot},\text{cop}}$, with $\gamma \in \pi$.

Theorem 5.3. $D(H)$ is a T-coalgebra. Moreover, the multiplication in $D(H)$ is uniquely defined by (13) and the condition that the embeddings $\overline{H}, H^{*\text{tot},\text{cop}} \hookrightarrow D(H)$ are T-coalgebra morphisms.

Proof. Firstly, for any $\alpha \in \pi$, we will show that $D_\alpha(H)$ is an associative algebra with unit. Then we will show that Δ , defined as above, is multiplicative, i.e., that any $\Delta_{\alpha,\beta}$ is an algebra morphism. After that, we will show that ε is an algebra morphism. Finally, we will check axioms for the antipode and that the conjugation isomorphisms are compatible with the multiplication.

ASSOCIATIVITY. Let α be in π . The multiplication defined in (12) is associative if and only if, for any $h, k, l \in H_{\alpha^{-1}}$, $p \in H_{\beta}^*$, $q \in H_{\gamma}^*$, and $r \in H_{\delta}^*$, with $\beta, \gamma, \delta \in \pi$,

$$(17) \quad ((h \otimes p)(k \otimes q))(l \otimes r) = (h \otimes p)((k \otimes q)(l \otimes r))$$

By computing the left-hand side of (17), we obtain

$$\begin{aligned} ((h \otimes p)(k \otimes q))(l \otimes r) &= h''_{(\alpha^{-1})} k''_{(\alpha^{-1})} l \otimes p \left\langle q, s_{\gamma}^{-1}(h''''_{(\gamma^{-1})}) \lrcorner \varphi_{\alpha}(h'_{(\alpha^{-1}\gamma\alpha)}) \right\rangle \\ &\quad \left\langle r, s_{\delta}^{-1}(h''''_{(\delta^{-1})} k''''_{(\delta^{-1})}) \lrcorner \varphi_{\alpha}(h''_{(\alpha^{-1}\delta\alpha)} k'_{(\alpha^{-1}\delta\alpha)}) \right\rangle \end{aligned}$$

(by the antimultiplicativity of s and the multiplicativity of φ)

$$\begin{aligned} &= h''_{(\alpha^{-1})} k''_{(\alpha^{-1})} l \otimes p \left\langle q, s_{\gamma}^{-1}(h''''_{(\gamma^{-1})}) \lrcorner \varphi_{\alpha}(h'_{(\alpha^{-1}\gamma\alpha)}) \right\rangle \\ &\quad \left\langle r, s_{\delta}^{-1}(k''''_{(\delta^{-1})}) s_{\delta}^{-1}(h''''_{(\delta^{-1})}) \lrcorner \varphi_{\alpha}(h''_{(\alpha^{-1}\delta\alpha)}) \varphi_{\alpha}(k'_{(\alpha^{-1}\delta\alpha)}) \right\rangle, \end{aligned}$$

while, by computing the right-hand side, we obtain

$$\begin{aligned} (h \otimes p)((k \otimes q)(l \otimes r)) &= h''_{(\alpha^{-1})} k''_{(\alpha^{-1})} l \otimes p \left\langle q \left\langle r, s_{\delta}^{-1}(k''''_{(\delta^{-1})}) \lrcorner \varphi_{\alpha}(k'_{(\alpha^{-1}\delta\alpha)}) \right\rangle, \right. \\ &\quad \left. s_{\gamma\delta}^{-1}(h''''_{(\delta^{-1}\gamma^{-1})}) \lrcorner \varphi_{\alpha}(h'_{(\alpha^{-1}\gamma\delta\alpha)}) \right\rangle \end{aligned}$$

(by the anticomultiplicativity of s and the comultiplicativity of φ)

$$\begin{aligned} &= h''_{(\alpha^{-1})} k''_{(\alpha^{-1})} l \otimes p \left\langle q, s_{\gamma}^{-1}(h''''_{(\gamma^{-1})}) \lrcorner \varphi_{\alpha}(h'_{(\alpha^{-1}\gamma\alpha)}) \right\rangle \\ &\quad \left\langle r, s_{\delta}^{-1}(k''''_{(\delta^{-1})}) s_{\delta}^{-1}(h''''_{(\delta^{-1})}) \lrcorner \varphi_{\alpha}(h''_{(\alpha^{-1}\delta\alpha)}) \varphi_{\alpha}(k'_{(\alpha^{-1}\delta\alpha)}) \right\rangle. \end{aligned}$$

UNIT. Let α be in π . For any $h \in H_{\alpha^{-1}}$ and $f \in H_{\gamma}^*$, with $\gamma \in \pi$, we have

$$(1_{\alpha^{-1}} \otimes \varepsilon)(h \otimes f) = 1_{\alpha^{-1}} h \otimes \varepsilon f = h \otimes f$$

and

$$(h \otimes f)(1_{\alpha} \otimes \varepsilon) = h''_{(\alpha^{-1})} 1_{\alpha^{-1}} \otimes f \varepsilon \langle \varepsilon, s_1^{-1}(h''') \rangle \langle \varepsilon, \varphi_{\alpha}(h') \rangle = h \otimes f,$$

where the fact that both s_1 and φ_{α} commute with ε .

MULTIPLICATIVITY OF Δ . Let us prove that $\Delta_{\alpha,\beta}$ is an algebra morphism for any $\alpha, \beta \in \pi$. Since $\Delta_{\alpha,\beta}$ obviously preserves the unit, we only need to prove that, for any $h, k \in H_{\beta^{-1}\alpha^{-1}}$, $f \in H_{\gamma}^*$ and $g \in H_{\delta}^*$, with $\gamma, \delta \in \pi$, we have

$$(18) \quad \Delta_{\alpha,\beta}((h \otimes f)(k \otimes g)) = \Delta_{\alpha,\beta}(h \otimes f) \Delta_{\alpha,\beta}(k \otimes g).$$

This is proved by evaluating both terms in (18) against the general term $p \otimes x \otimes q \otimes y$ ($p \in H_{\alpha^{-1}}^*$, $q \in H_{\beta^{-1}}^*$, and $x, y \in H_{\gamma\delta}$).

MULTIPLICATIVITY OF ε . For any $h, k \in H_1$, $f \in H_{\gamma}^*$, and $g \in H_{\delta}^*$, with $\gamma, \delta \in \pi$, we have

$$\langle \varepsilon, h \otimes f \rangle \langle \varepsilon, k \otimes g \rangle = \langle \varepsilon, h \rangle \langle f, 1_{\gamma} \rangle \langle \varepsilon, k \rangle \langle g, 1_{\delta} \rangle$$

and

$$\begin{aligned} \langle \varepsilon, (h \otimes f)(k \otimes g) \rangle &= \left\langle \varepsilon, h'' k \otimes f \left\langle g, s_{\delta}^{-1}(h''_{(\delta^{-1})}) \lrcorner h'_{(\delta)} \right\rangle \right\rangle \\ &= \langle \varepsilon, h'' \rangle \langle \varepsilon, k \rangle \langle f, 1_{\gamma} \rangle \langle g, s_{\delta}^{-1}(h''_{(\delta^{-1})}) h'_{(\delta)} \rangle \\ &= \langle \varepsilon, k \rangle \langle f, 1_{\gamma} \rangle \langle g, s_{\delta}^{-1}(h''_{(\delta^{-1})}) h'_{(\delta)} \rangle \\ &= \langle \varepsilon, k \rangle \langle f, 1_{\gamma} \rangle \langle g, \langle \varepsilon, h \rangle 1_{\delta} \rangle = \langle \varepsilon, h \rangle \langle f, 1_{\gamma} \rangle \langle \varepsilon, k \rangle \langle g, 1_{\delta} \rangle. \end{aligned}$$

This proves that ε is multiplicative. Moreover, since ε is obviously unitary, it is an algebra homomorphism.

ANTIPODE. Let h be in H_1 and let f be in H_γ^* , with $\gamma \in \pi$. We have

$$\begin{aligned} & (h \otimes f)'_{(\alpha)} s_{\alpha^{-1}}((h \otimes f)''_{(\alpha^{-1})}) \\ &= (\varphi_{\alpha^{-1}}(h'_{(\alpha^{-1})}) \otimes f')((\varphi_{\alpha^{-1}} \circ s_\alpha)(h''_{(\alpha)}) \otimes \varepsilon)(1_{\alpha^{-1}} \otimes s_*(f'')) \\ &= (\varphi_{\alpha^{-1}}(h'_{(\alpha^{-1})} s_\alpha(h''_{(\alpha)})) \otimes f')(1_{\alpha^{-1}} \otimes s_*(f'')) \\ &= \langle \varepsilon, h \rangle (1_{\alpha^{-1}} \otimes f')(1 \otimes s_*(f'')) = \langle \varepsilon, h \rangle 1 \otimes f' s_*(f'') \\ &= \langle \varepsilon, h \rangle \langle f, 1_\gamma \rangle 1_{\alpha^{-1}} \otimes \varepsilon = \langle \varepsilon, h \otimes f \rangle 1_{\alpha^{-1}} \otimes \varepsilon \end{aligned}$$

and

$$\begin{aligned} & s_{\alpha^{-1}}((h \otimes f)'_{(\alpha^{-1})})(h \otimes f)''_{(\alpha)} = s_{\alpha^{-1}}(\varphi_\alpha(h'_{(\alpha)}) \otimes f')(h''_{(\alpha^{-1})} \otimes f'') \\ &= (s_\alpha(h'_\alpha) \otimes \varepsilon)(1_\alpha \otimes s_*(f'))(h''_{(\alpha^{-1})} \otimes f'') \\ &= \langle f, 1_\gamma \rangle (s_\alpha(h'_\alpha) \otimes \varepsilon)(h''_{(\alpha^{-1})} \otimes \varepsilon) = \langle f, 1_\gamma \rangle s_\alpha(h''_{(\alpha)}) h'''_{(\alpha^{-1})} \otimes \langle \varepsilon, h' \rangle \langle \varepsilon, h'' \rangle \varepsilon \\ &= \langle \varepsilon, h \rangle \langle f, 1_\gamma \rangle 1_\alpha \otimes \varepsilon = \langle \varepsilon, h \otimes f \rangle 1_\alpha \otimes \varepsilon. \end{aligned}$$

CONJUGATION. Let us check that φ_β^α is an algebra isomorphism for any $\alpha, \beta \in \pi$. Since φ_β^α is obviously bijective and preserve the unit, we only need to show that it is compatible with the multiplication, i.e., that, for any $h, k \in H_{\alpha^{-1}}$, $f \in H_\gamma^*$, and $g \in H_\delta^*$, with $\gamma, \delta \in \pi$, we have

$$(19) \quad \varphi_\beta(h \otimes f) \varphi_\beta(k \otimes g) = \varphi_\beta((h \otimes f)(k \otimes g)).$$

This is proved by evaluating both sides in (19) against the general term $p \otimes x$ ($x \in H_{\beta\gamma\delta\beta^{-1}}$ and $p \in H_{\beta\alpha^{-1}\beta^{-1}}^*$).

This concludes the proof that $D(H)$ is a T-coalgebra.

EMBEDDINGS. It only remains to show that (13), together with the request that the canonical embeddings $\overline{H}, H^{*\text{tot}, \text{cop}} \hookrightarrow D(H)$ are T-coalgebra morphisms, uniquely determinates the multiplication in $D(H)$. Let α be in π . For any $h, k \in H_{\alpha^{-1}}$, $f \in H_\gamma^*$ and $g \in H_\delta^*$ we have

$$(h \otimes f)(k \otimes g) =$$

(by (13a))

$$= (1_{\alpha^{-1}} \otimes f)(h \otimes \varepsilon)(1_{\alpha^{-1}} \otimes g)(k \otimes \varepsilon)$$

(by (13b))

$$= (1_{\alpha^{-1}} \otimes f) \left(h''_{(\alpha^{-1})} \otimes \langle g, s_\delta^{-1}(h'''_{(\delta^{-1})}) - \varphi_\alpha(h'_{(\alpha^{-1}\delta\alpha)}) \rangle \right) (k \otimes \varepsilon)$$

(by (13a))

$$= (1_{\alpha^{-1}} \otimes f) \left(1_{\alpha^{-1}} \otimes \langle g, s_\delta^{-1}(h'''_{(\delta^{-1})}) - \varphi_\alpha(h'_{(\alpha^{-1}\delta\alpha)}) \rangle \right) (h''_{(\alpha^{-1})} \otimes \varepsilon)(k \otimes \varepsilon)$$

(since the canonical embedding of both $H_{\alpha^{-1}}$ and $H_\alpha^{*\text{tot}, \text{cop}}$ in $D_\alpha(H)$ are algebra morphisms)

$$= \left(1_{\alpha^{-1}} \otimes f \langle g, s_\delta^{-1}(h'''_{(\delta^{-1})}) - \varphi_\alpha(h'_{(\alpha^{-1}\delta\alpha)}) \rangle \right) (h''_{(\alpha^{-1})} k \otimes \varepsilon)$$

(again by (13a))

$$= h''_{(\alpha^{-1})} k \otimes f \left\langle g, s_\delta^{-1}(h'''_{(\delta^{-1})}) - \varphi_\alpha(h'_{(\alpha^{-1}\delta\alpha)}) \right\rangle.$$

This concludes the proof of the theorem.

Coassociativity, compatibility between the comultiplication and the counit, compatibility between the comultiplication and the conjugation, and, finally, the fact that φ is a group homomorphism follow by observing that $D(H)$ has the same comultiplication, counit, and The omitted computations are the same needed conjugation of the tensor product of T-coalgebras $\overline{H} \otimes H^{*\text{tot}, \text{cop}}$. $\square$

QUASITRIANGULAR STRUCTURE OF THE QUANTUM DOUBLE. To prove that $D(H)$ is quasitriangular, we need some preliminary results. For any $\alpha \in \pi$, we set $n_\alpha = \dim H_\alpha$. Let $(e_{\alpha,i})_{i=1,\dots,n_\alpha}$ be a basis of H_α as a vector space and let $(e^{\alpha,i})_{i=1,\dots,n_\alpha}$ be the dual basis of $(e_{\alpha,i})_{i=1,\dots,n_\alpha}$. We set

$$(20) \quad R_{\alpha,\beta} = e_{\alpha^{-1},i} \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes e^{\alpha^{-1},i} \in D_\alpha(H) \otimes D_\beta(H)$$

$$\text{and } \tilde{R}_{\alpha,\beta} = s_\alpha(e_{\alpha,i}) \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes e^{\alpha,i}.$$

Lemma 5.4. *For any $\alpha, \beta \in \pi$, both $R_{\alpha,\beta}$ and $\tilde{R}_{\alpha,\beta}$ are independent of the choice of the bases. Moreover, $\tilde{R}_{\alpha,\beta}$ is the inverse of $R_{\alpha,\beta}$ in the algebra $D_\alpha(H) \otimes D_\beta(H)$.*

Proof. Let $\mathfrak{C}_{\alpha,\beta}(H)$ be the subspace of $D_\alpha(H) \otimes D_\beta(H)$ generated by the elements $h \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes F$ with $h \in H_{\alpha^{-1}}$ and $F \in H_\alpha^{*\text{tot}, \text{cop}}$. Clearly, $\mathfrak{C}_{\alpha,\beta}(H)$ is a subalgebra of $D_\alpha(H) \otimes D_\beta(H)$. Moreover, we have an equivalence of algebras $\text{Conv}_{\alpha^{-1}}(H) \rightarrow \mathfrak{C}_{\alpha,\beta}(H)$: $h \otimes F \mapsto (h \otimes \varepsilon) \otimes (1_{\beta^{-1}} \otimes F)$, for any $h \in H_{\alpha^{-1}}$ and $F \in H_\alpha^{*\text{tot}, \text{cop}}$. In particular, $R_{\alpha,\beta}$ is the image of the identity morphism $e_{\alpha^{-1},i} \otimes e^{\alpha^{-1},i}$ of $H_{\alpha^{-1}}$ under this isomorphism. Since the α -th component of the antipode s of H is the inverse, in the algebra $\text{Conv}_{\alpha^{-1}}(H)$, of $e_{\alpha^{-1},i} \otimes e^{\alpha^{-1},i}$, also $R_{\alpha,\beta}$ is invertible. Since s_α can be represented as $s(e_{\alpha,i}) \otimes e^{\alpha,i}$ and $\tilde{R}_{\alpha,\beta}$ is the image of s_α under the above isomorphism, we conclude that $\tilde{R}_{\alpha,\beta}$ is the inverse of $R_{\alpha,\beta}$. $\square$

Remark 5.5. Let h be in H_α , with $\alpha \in \pi$, and suppose $\alpha_1 \alpha_2 \cdots \alpha_n = \alpha$ for certain $\alpha_1, \alpha_2, \dots, \alpha_n \in \pi$. By observing that $h = h^i e_{\alpha,i} = \langle e^{\alpha,i}, h \rangle e_{\alpha,i}$ and by the linearity of Δ we have

$$(21) \quad h'_{(\alpha_1)} \otimes h''_{(\alpha_2)} \cdots \otimes h'^n_{(\alpha_n)} = \langle e^{\alpha,i}, h \rangle (e_{\alpha,i})'_{(\alpha_1)} \otimes (e_{\alpha,i})''_{(\alpha_2)} \otimes \cdots \otimes (e_{\alpha,i})'^n_{(\alpha_n)}.$$

Theorem 5.6. *$D(H)$ is quasitriangular with universal R -matrix given by (20) for any $\alpha, \beta \in \pi$.*

Proof. $R_{\alpha,\beta}$ is well defined and invertible by Lemma 5.4. We still need to check the four relations (7).

RELATION (7a). Let α, β , and γ be π . Given $h \in H_{\beta^{-1}\alpha^{-1}}$ and $f \in H_\gamma^*$, we have

$$\begin{aligned} R_{\alpha,\beta} \Delta_{\alpha,\beta}(h \otimes f) &= ((e_{\alpha^{-1},i} \otimes \varepsilon) \otimes (1_{\beta^{-1}} \otimes e^{\alpha^{-1},i})) ((\varphi_\beta(h'_{(\beta^{-1}\alpha^{-1}\beta)}) \otimes f') \otimes (h''_{(\beta^{-1})} \otimes f'')) \\ &= (e_{\alpha^{-1},i})''_{(\alpha^{-1})} \varphi_\beta(h'_{(\beta^{-1}\alpha\beta)}) \otimes \\ &\quad \otimes \left\langle f', s_\gamma^{-1}((e_{\alpha^{-1},i})'''_{(\gamma^{-1})}) - \varphi_\alpha((e_{\alpha^{-1},i})'_{(\alpha^{-1}\gamma\alpha)}) \right\rangle \otimes h''_{(\beta^{-1})} \otimes e^{\alpha^{-1},i} f''. \end{aligned}$$

Since the action of $(\sigma \circ (\varphi_\alpha \otimes D_\alpha(H)) \circ \Delta_{\alpha\beta\alpha^{-1},\alpha})(-)$ on $h \otimes f$ is given by

$$\begin{aligned} h \otimes f &\xrightarrow{\Delta_{\alpha\beta\alpha^{-1},\alpha}} \varphi_\alpha(h'_{(\beta^{-1})}) \otimes f' \otimes h''_{(\alpha^{-1})} \otimes f'' \xrightarrow{\varphi_{\alpha^{-1}} \otimes D_\alpha(H)} \\ &h'_{(\beta^{-1})} \otimes \varphi_\alpha^*(f') \otimes h''_{(\alpha^{-1})} \otimes f'' \xrightarrow{\sigma} h''_{(\alpha^{-1})} \otimes f'' \otimes h'_{(\beta^{-1})} \otimes \varphi_\alpha^*(f'), \end{aligned}$$

we have

$$\begin{aligned}
& \left(\left(\sigma \circ (\varphi_\alpha \otimes D_\alpha(H)) \right) \circ \Delta_{\alpha\beta\alpha^{-1},\alpha} \right) (h \otimes f) \Big) R_{\alpha,\beta} \\
&= (h''_{(\alpha^{-1})} \otimes f'' \otimes h'_{(\beta^{-1})} \otimes \varphi_\alpha^*(f')) (e_{\alpha^{-1},i} \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes e^{\alpha^{-1},i}) \\
&= h'''_{(\alpha^{-1})} e_{\alpha^{-1},i} \otimes f'' \otimes h''_{(\beta^{-1})} \otimes \varphi_\alpha^*(f') \langle e^{\alpha^{-1},i}, s_{\alpha^{-1}}^{-1}(h'''_{(\alpha)})_{-\varphi_\beta}(h'_{(\beta^{-1}\alpha^{-1}\beta)}) \rangle.
\end{aligned}$$

Relation (7a) is proved by observing that evaluating the two expressions above against the tensor $H_{\alpha^{-1}} \otimes H_\gamma^* \otimes H_{\beta^{-1}} \otimes \langle \cdot, x \rangle$ (for a generic $x \in H_{\alpha^{-1}\gamma}$) we found the same result.

RELATION (7b). For any $\alpha, \beta, \gamma \in \pi$, we have

$$\begin{cases} (R_{\alpha,\gamma})_{1\beta 3} = (e_{\alpha^{-1},i} \otimes \varepsilon) \otimes (1_{\beta^{-1}} \otimes \varepsilon) \otimes (1_{\gamma^{-1}} \otimes e^{\alpha^{-1},i}) \\ (R_{\alpha,\beta})_{12\gamma} = (e_{\alpha^{-1},i} \otimes \varepsilon) \otimes (1_{\beta^{-1}} \otimes e^{\alpha^{-1},i}) \otimes (1_{\gamma^{-1}} \otimes \varepsilon) \end{cases}$$

and so we have $(R_{\alpha,\gamma})_{1\beta 3} (R_{\alpha,\beta})_{12\gamma} = e_{\alpha^{-1},i} e_{\alpha^{-1},j} \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes e^{\alpha^{-1},j} \otimes 1_{\gamma^{-1}} \otimes e^{\alpha^{-1},i}$. Since $R_{\alpha,\beta\gamma} = e_{\alpha^{-1},i} \otimes \varepsilon \otimes 1_{(\beta\gamma)^{-1}} \otimes e^{\alpha^{-1},i}$, we have $(D_\alpha(H) \otimes \Delta_{\beta,\gamma})(R_{\alpha,\beta\gamma}) = e_{\alpha^{-1},i} \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes (e^{\alpha^{-1},i})' \otimes 1_{\gamma^{-1}} \otimes (e^{\alpha^{-1},i})''$. So, we only need to prove

$$(22) \quad e_{\alpha^{-1},i} e_{\alpha^{-1},j} \otimes e^{\alpha^{-1},j} \otimes e^{\alpha^{-1},i} = e_{\alpha^{-1},i} \otimes (e^{\alpha^{-1},i})' \otimes (e^{\alpha^{-1},i})''.$$

If we evaluate both sides of (22) against the tensor $f \otimes H_{\alpha^{-1}}^* \otimes H_{\alpha^{-1}}^*$ (for a generic $f \in H_\alpha^*$), then, by (21), we found the same result.

RELATION (7c). Let α, β , and γ be in π . Observing that $R_{\beta^{-1}\alpha\beta,\gamma} = e_{\beta^{-1}\alpha^{-1}\beta,i} \otimes \varepsilon \otimes 1_{\gamma^{-1}} \otimes e^{\beta^{-1}\alpha^{-1}\beta,i}$, we obtain

$$\left((\varphi_\beta \otimes D_\gamma(H)) (R_{\beta^{-1}\alpha\beta,\gamma}) \right)_{1\beta 3} = \varphi_\beta(e_{\beta^{-1}\alpha^{-1}\beta,i}) \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes \varepsilon \otimes 1_{\gamma^{-1}} \otimes e^{\beta^{-1}\alpha^{-1}\beta,i}.$$

Observing that $(R_{\beta,\gamma})_{\alpha 23} = 1_{\alpha^{-1}} \otimes \varepsilon \otimes e_{\beta^{-1},i} \otimes \varepsilon \otimes 1_{\gamma^{-1}} \otimes e^{\beta^{-1},i}$, we obtain $\left((\varphi_\beta \otimes D(H)_\gamma) (R_{\beta^{-1}\alpha\beta,\gamma}) \right)_{1\beta 3} (R_{\beta,\gamma})_{\alpha 23} = \varphi_\beta(e_{\beta^{-1}\alpha^{-1}\beta,i}) \otimes \varepsilon \otimes e_{\beta^{-1},j} \otimes \varepsilon \otimes 1_{\gamma^{-1}} \otimes e^{\beta^{-1}\alpha^{-1}\beta,i} e^{\beta^{-1},j}$. Finally, observing that $R_{\alpha\beta,\gamma} = e_{(\alpha\beta)^{-1},i} \otimes \varepsilon \otimes 1_{\gamma^{-1}} \otimes e^{(\alpha\beta)^{-1},i}$, we obtain $(\Delta_{\alpha,\beta} \otimes D_\gamma(H)) (R_{\alpha\beta,\gamma}) = \varphi_\beta((e_{(\alpha\beta)^{-1},i})'_{(\beta^{-1}\alpha^{-1}\beta)}) \otimes \varepsilon \otimes (e_{(\alpha\beta)^{-1},i})''_{(\beta^{-1})} \otimes \varepsilon \otimes 1_{\gamma^{-1}} \otimes e^{(\alpha\beta)^{-1},i}$. So, to prove (7c), we only need to show the equality

$$\begin{aligned}
(23) \quad & e_{\beta^{-1}\alpha^{-1}\beta,i} \otimes e_{\beta^{-1},j} \otimes e^{\beta^{-1}\alpha^{-1}\beta,i} e^{\beta^{-1},j} \\
&= (e_{(\alpha\beta)^{-1},i})'_{(\beta^{-1}\alpha\beta)} \otimes (e_{(\alpha\beta)^{-1},i})''_{(\beta^{-1})} \otimes e^{(\alpha\beta)^{-1},i}.
\end{aligned}$$

If we evaluate both sides of (23) against the tensor $H_{\beta^{-1}\alpha^{-1}\beta} \otimes H_{\beta^{-1}} \otimes \langle \cdot, x \rangle$ (for a generic $x \in H_{\beta^{-1}\alpha^{-1}}$), then by (21) we found the same result.

RELATION (7d). Given $\alpha, \beta, \gamma \in \pi$, we have $(\varphi_\gamma \otimes \varphi_\gamma)(R_{\alpha,\beta}) = (\varphi_\gamma \otimes \varphi_\gamma)(e_{\alpha^{-1},i} \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes e^{\alpha^{-1},i}) = \varphi_\gamma(e_{\alpha^{-1},i}) \otimes \varepsilon \otimes 1_{\beta^{-1}} \otimes \varphi_\gamma(e^{\alpha^{-1},i})$. Now, φ_γ is a linear isomorphism, so $(\varphi_\gamma(e_{\alpha^{-1},i}))_{i=1,\dots,n_\alpha}$ is a basis of $H_{\gamma\alpha^{-1}\gamma^{-1}}$, and $(\varphi_\gamma^*(e^{\alpha^{-1},i}))_{i=1,\dots,n_\alpha}$ is its dual basis. So, by Lemma 5.4, $R_{\gamma\alpha\gamma^{-1},\gamma\beta\gamma^{-1}} = (\varphi_\gamma \otimes \varphi_\gamma)(R_{\alpha,\beta})$.

This concludes the proof of the theorem. $\square$

PROOF OF THEOREM 5.1. We need a preliminary lemma.

Lemma 5.7. *Let T be any quasitriangular T-coalgebra with R-matrix R . For any $\alpha, \beta \in \pi$ and any $x \in T_\alpha$ we have*

$$(24) \quad s_\beta^{-1}(x'''_{(\beta^{-1})}) \xi_{(\beta),i} x'_{(\beta)} \otimes \zeta_{(\beta^{-1}\alpha\beta)} x''_{(\beta^{-1}\alpha\beta)} = \xi_{(\beta),i} \otimes \varphi_{\beta^{-1}}(x) \zeta_{(\beta^{-1}\alpha\beta)}.$$

Proof. By (7a), we have $\xi_{(\beta).i}x'_{(\beta)} \otimes \zeta_{(\beta^{-1}\alpha\beta).i}x''_{(\beta^{-1}\alpha\beta)} \otimes x'''_{(\beta^{-1})} = x''_{(\beta^{-1})}\xi_{(\beta).i} \otimes \varphi_{\beta^{-1}}(x'_{(\alpha)})\zeta_{(\beta^{-1}\alpha\beta).i} \otimes x'''_{(\beta^{-1})}$. If, on both sides, we apply $T_\beta \otimes T_{\beta^{-1}\alpha\beta} \otimes s_\beta^{-1}$ and exchange the first and the third factors, then we obtain $s_\beta^{-1}(x'''_{(\beta^{-1})}) \otimes \xi_{(\beta).i}x'_{(\beta)} \otimes \zeta_{(\beta^{-1}\alpha\beta).i}x''_{(\beta^{-1}\alpha\beta)} = s_\beta^{-1}(x'''_{(\beta^{-1})}) \otimes x''_{(\beta).i}\xi_{(\beta).i} \otimes \varphi_{\beta^{-1}}(x'_{(\alpha)})\zeta_{(\beta^{-1}\alpha\beta).i}$. Applying $\mu_\beta \otimes T_{\beta^{-1}\alpha\beta}$ on both sides, and observing that $s_\beta^{-1}(x'''_{(\beta^{-1})})x''_{(\beta)} \otimes \varphi_{\beta^{-1}}(x'_{(\alpha)}) = \varepsilon(x'_{(\beta)})1_\beta \otimes \varphi_{\beta^{-1}}(x'_{(\alpha)}) = 1_\beta \otimes \varphi_{\beta^{-1}}(x)$, we obtain (24). $\square$

Proof of Theorem 5.1. By Theorem 5.6, the T-coalgebra $D(H)$ defined as in Theorem 5.3 satisfies the three conditions of Theorem 5.1. We still have to check that these three conditions determinate the T-coalgebra structure on $D(H)$, i.e., that the multiplication and the comultiplication we gave in the definition of $D(H)$ are uniquely defined by the requirements of Theorem 5.1.

PRODUCT. Suppose that $D(H)$ satisfies the three conditions in Theorem 5.1. Let us check, that, given $\alpha \in \pi$, the multiplication on $D_\alpha = H_{\alpha^{-1}} \otimes H^{*\text{tot},\text{cop}}$ must satisfy the relations (13). Let p_α be defined as in (11). The bijectivity of p_α implies (13a). Let h be in $\overline{H}_\alpha$ and let $f \in H_\gamma^* \subseteq H_1^{*\text{tot},\text{cop}}$, with $\gamma \in \pi$. By Lemma 24, we have

$$(25) \quad s_{\gamma^{-1}}^{-1}((h \otimes \varepsilon)'''_{(\gamma)}) (e_{\gamma.i} \otimes \varepsilon) (h \otimes \varepsilon)'_{(\gamma^{-1})} \otimes (1_{\gamma\alpha^{-1}\gamma^{-1}} \otimes e^{\gamma.i}) (h \otimes \varepsilon)''_{(\gamma\alpha\gamma^{-1})} \\ = (e_{\gamma.i} \otimes \varepsilon) \otimes \varphi_\gamma(h \otimes \varepsilon) (1_{\gamma\alpha\gamma^{-1}} \otimes e^{\gamma.i}).$$

By observing that $h'_{(\gamma^{-1})} \otimes h''_{(\gamma\alpha\gamma^{-1})} \otimes h'''_{(\gamma)} = \varphi_\gamma(h'_{(\alpha^{-1}\gamma\alpha)}) \otimes \varphi_\gamma(h'_{(\alpha^{-1})}) \otimes h'''_{\gamma^{-1}}$, if we compute the left-hand side in (25), then we obtain

$$s_{\gamma^{-1}}^{-1}((h \otimes \varepsilon)'''_{(\gamma)}) (e_{\gamma.i} \otimes \varepsilon) (h \otimes \varepsilon)'_{(\gamma^{-1})} \otimes (1_{\gamma\alpha^{-1}\gamma^{-1}} \otimes e^{\gamma.i}) (h \otimes \varepsilon)''_{(\gamma\alpha\gamma^{-1})} \\ = ((\varphi_{\gamma^{-1}} \circ s_\gamma)^{-1}(h'''_{(\gamma^{-1})}) \otimes \varepsilon) (e_{\gamma.i} \otimes \varepsilon) (\varphi_\gamma(h'_{(\alpha^{-1}\gamma\alpha)}) \otimes \varepsilon) \otimes \\ \otimes (1_{\gamma\alpha^{-1}\gamma^{-1}} \otimes e^{\gamma.i}) (\varphi_\gamma(h'_{(\alpha^{-1})}) \otimes \varepsilon) \\ = ((\varphi_\gamma \circ s_\gamma^{-1})(h'''_{(\gamma^{-1})})e_{\gamma.i}\varphi_\gamma(h'_{(\alpha^{-1}\gamma\alpha)}) \otimes \varepsilon) \otimes (\varphi_\gamma(h'_{(\alpha^{-1})}) \otimes e^{\gamma.i}),$$

where, in the last passage, we used both the fact that the immersion of $\overline{H}$ in $D(H)$ is a morphism of T-coalgebras and (13a). So, if we apply $\varphi_{\gamma^{-1}} \otimes \varphi_{\gamma^{-1}}$ on both sides of (25), then we get

$$(26) \quad (s_\gamma^{-1}(h'''_{(\gamma^{-1})})e_{\gamma.i}\varphi_\alpha(h'_{(\alpha^{-1}\gamma\alpha)}) \otimes \varepsilon) \otimes (h''_{(\alpha)} \otimes e^{\gamma.i}) \\ = (e_{\gamma.i} \otimes \varepsilon) \otimes (h \otimes \varepsilon)(1_{\alpha^{-1}} \otimes e^{\gamma.i}).$$

If we evaluate both terms of (26) against $\langle f \otimes 1_1, _ \rangle \otimes D_\alpha(H)$, then on the left-hand side we get $\langle f, s_\gamma^{-1}e_{\gamma.i}\varphi_\alpha(h'_{(\alpha^{-1}\gamma\alpha)}) \rangle h''_{(\alpha)} \otimes e^{\gamma.i} = h''_{(\alpha)} \otimes \langle f, s_\gamma^{-1}\varphi_\alpha(h'_{(\alpha^{-1}\gamma\alpha)}) \rangle$, while on the right-hand side we get $\langle f, e_{\gamma.i} \rangle (h \otimes \varepsilon) (1_{\alpha^{-1}} \otimes e^{\gamma.i}) = (h \otimes \varepsilon) (1_{\alpha^{-1}} \otimes f)$.

COMULTIPLICATION. Let us check that the comultiplication on $D(H)$ is also unique. Given $\alpha, \beta, \gamma \in \pi$, $h \in \overline{H}_{\alpha\beta}$, and $f \in H_\gamma^*$, we have

$$\Delta_{\alpha,\beta}(h \otimes f) = \Delta_{\alpha,\beta}((1_{\alpha^{-1}} \otimes f) (h \otimes \varepsilon)) \\ = (1_{\alpha^{-1}} \otimes f') (h'_{(\alpha)} \otimes \varepsilon) \otimes (1_{\alpha^{-1}} \otimes f'') (h''_{(\beta)} \otimes \varepsilon) \\ = (\varphi_\beta(h'_{(\beta^{-1}\alpha^{-1}\beta)}) \otimes f') \otimes (h''_{(\beta^{-1})} \otimes f'').$$

$\square$

Remark 5.8. The quantum double of a Hopf algebra can be obtained also via the Majid bicrossproduct [9]. This is true also in the crossed case [25]. Recently,

Vainerman [21] constructed some examples of nontrivial T-coalgebras with non-isomorphic components. These examples can also be interpreted —and, eventually, enlarged— as bicrossproducts of T-coalgebras.

Remark 5.9. Street [16] has proved that, starting from any (not necessarily finite-dimensional) Hopf algebra H , it is possible to construct, via Tannaka Theory, a coquasitriangular Hopf algebra $D^*(H)$ such that, when H is finite-dimensional, $D^*(H) = (D(H))^*$. Tannaka Theory for T-algebras was developed by the author in [26] where, contextually, it is provided an analog for the co-double construction [12] in the case of a T-algebra.

6. RIBBON T-COALGEBRAS

The notion of a ribbon Hopf algebra [13] is generalized to the case of a T-coalgebra in [20].

DRINFELD ELEMENTS. Following [22], we set $u_\alpha = (s_{\alpha^{-1}} \circ \varphi_\alpha)(\zeta_{(\alpha^{-1}).i})\xi_{(\alpha).i}$ and $u = \{u_\alpha\}_{\alpha \in \pi}$. The u_α are called *Drinfeld elements* of H . When $\pi = \{1\}$, we recover the usual definition of Drinfeld element of a quasitriangular Hopf algebra.

The following properties of u are proved in [22]. Let α and β be in π and let h be in H_α .

- (27a) $u_1 = s_1(\zeta_{(1).i})\xi_{(1).i}$.
- (27b) u_α is invertible with inverse $u_\alpha^{-1} = s_\alpha^{-1}(\tilde{\zeta}_{(\alpha^{-1}).i})\tilde{\xi}_{(\alpha).i}$. Moreover we have $u_\alpha^{-1} = (s_\alpha^{-1} \circ s_{\alpha^{-1}})(\zeta_{(\alpha).i})\xi_{(\alpha).i} = \xi_{(\alpha).i}(s_{\alpha^{-1}} \circ s_\alpha)(\zeta_{(\alpha).i})$.
- (27c) $(u_{\alpha\beta})'_{(\alpha)} \otimes (u_{\alpha\beta})''_{(\beta)} = \tilde{\xi}_{(\alpha).i}\tilde{\zeta}_{(\alpha).j}u_\alpha \otimes \tilde{\zeta}_{(\beta).i}\varphi_{\alpha^{-1}}(\tilde{\xi}_{(\alpha\beta\alpha^{-1}).j})u_\beta$.
- (27d) $\varepsilon(u_1) = 1$.
- (27e) $s_{\alpha^{-1}}(u_{\alpha^{-1}})u_\alpha = u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}})$.
- (27f) $\varphi_\beta(u_\alpha) = u_{\beta\alpha\beta^{-1}}$.
- (27g) $(s_{\alpha^{-1}} \circ s_\alpha \circ \varphi_\alpha)(h) = u_\alpha h u_\alpha^{-1}$.
- (27h) $u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}})h = \varphi_{\alpha^2}(h)u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}})$.

Notice that, by (27g), we have $(s_{\alpha^{-1}} \circ s_\alpha)(h) = \varphi_{\alpha^{-1}}(u_\alpha)\varphi_{\alpha^{-1}}(h)\varphi_{\alpha^{-1}}(u_\alpha^{-1}) = u_\alpha\varphi_{\alpha^{-1}}(h)u_\alpha^{-1}$. In particular, for $h = u_\alpha$ we obtain

$$(27i) \quad (s_{\alpha^{-1}} \circ s_\alpha)(u_\alpha) = u_\alpha.$$

DEFINITION OF A RIBBON T-COALGEBRA (FIRST VERSION). Let H be a quasitriangular T-coalgebra. According to [20], we say that H is a *ribbon T-coalgebra* if it is endowed with a family $\theta = \{\theta_\alpha | \theta_\alpha \in H_\alpha\}_{\alpha \in \pi}$, the *twist*, such that θ_α is invertible for any $\alpha \in \pi$ and the following conditions are satisfied for any $\alpha, \beta \in \pi$ and $h \in H_\alpha$.

- (28a) $\varphi_\alpha(h) = \theta_\alpha^{-1}h\theta_\alpha$.
- (28b) $s_\alpha(\theta_\alpha) = \theta_{\alpha^{-1}}$.
- (28c) $(\theta_{\alpha\beta})'_{(\alpha)} \otimes (\theta_{\alpha\beta})''_{(\beta)} = \theta_\alpha\zeta_{(\alpha).i}\xi_{(\alpha).j} \otimes \theta_\beta\varphi_{\alpha^{-1}}(\xi_{(\alpha\beta\alpha^{-1}).i})\zeta_{(\beta).j}$.
- (28d) $\varphi_\beta(\theta_\alpha) = \theta_{\beta\alpha\beta^{-1}}$.

Notice that $(H_1, R_{1,1}, \theta_1)$ is a ribbon Hopf algebra in the usual sense.

If $H = (H, R, \theta)$ is a ribbon T-coalgebra, then, for any $\alpha \in \pi$, we obtain the following properties.

- (29a) $\varphi_{\alpha^{-1}}(h) = \theta_\alpha h \theta_\alpha^{-1}$ for any $h \in H_\alpha$ (this follows by 28a).
- (29b) $\varepsilon(\theta_1) = 1$ (this is because H_1 is a ribbon Hopf algebra).
- (29c) θ_1 is central (by the same reason).
- (29d) $\theta_\alpha\varphi_\alpha(h) = h\theta_\alpha$ for any $h \in H_\alpha$ (this follows by 28a).

Moreover, it is proved in [22] that we have

- (29e) $\theta_\alpha u_\alpha = u_\alpha \theta_\alpha$ and
- (29f) $\theta_\alpha^{-2} = s_{\alpha^{-1}}(u_{\alpha^{-1}})u_\alpha = u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}})$.

If for any $\alpha, \beta \in \pi$, we introduce the notation

$$(30a) \quad Q_{\alpha, \beta} = (\sigma \circ (\varphi_{\alpha^{-1}} \otimes H_{\alpha}))(R_{\alpha\beta\alpha^{-1}, \alpha})R_{\alpha, \beta} = \\ = \zeta_{(\alpha).i} \xi_{(\alpha).j} \otimes \varphi_{\alpha^{-1}}(\xi_{(\alpha\beta\alpha^{-1}).i}) \zeta_{(\beta).j},$$

then (28c) can be written in the form $(\theta_{\alpha\beta})'_{(\alpha)} \otimes (\theta_{\alpha\beta})''_{(\beta)} = (\theta_{\alpha} \otimes \theta_{\beta})Q_{\alpha, \beta}$. Moreover, if we set $\tilde{Q}_{\alpha, \beta} = Q_{\alpha, \beta}^{-1} = \tilde{\xi}_{(\alpha).i} \tilde{\zeta}_{(\alpha).j} \otimes \tilde{\zeta}_{(\beta).i} \varphi_{\alpha^{-1}}(\tilde{\xi}_{(\alpha\beta\alpha^{-1}).j})$, we can rewrite (27c) in the form

$$(30b) \quad (u_{\alpha\beta})'_{(\alpha)} \otimes (u_{\alpha\beta})''_{(\beta)} = \tilde{Q}_{\alpha, \beta}(u_{\alpha} \otimes u_{\beta}).$$

We observe that the conjugation preserves Q , i.e., that, for any $\alpha, \beta, \gamma \in \pi$, we have $(\varphi_{\alpha} \otimes \varphi_{\alpha})(Q_{\beta, \gamma}) = Q_{\alpha\beta\alpha^{-1}, \alpha\gamma\alpha^{-1}}$.

DEFINITION OF A RIBBON T-COALGEBRA (SECOND VERSION). We can define a ribbon T-coalgebra in an equivalent way as a quasitriangular T-coalgebra H endowed with a family $v = \{v_{\alpha} | v_{\alpha} \in H_{\alpha}\}_{\alpha \in \pi}$ satisfying the following conditions.

- (31a) $hv_{\alpha} = v_{\alpha}\varphi_{\alpha^{-1}}(h)$ for any $h \in H_{\alpha}$.
- (31b) $v_{\alpha}^2 = u_{\alpha}s_{\alpha^{-1}}(u_{\alpha^{-1}})$.
- (31c) $(v_{\alpha\beta})'_{(\alpha)} \otimes (v_{\alpha\beta})''_{(\beta)} = \tilde{Q}_{\alpha, \beta}(v_{\alpha} \otimes v_{\beta})$.
- (31d) $s_{\alpha}(v_{\alpha}) = v_{\alpha^{-1}}$.
- (31e) $\varphi_{\beta}(v_{\alpha}) = v_{\beta\alpha\beta^{-1}}$.

The proof of the equivalence of the two definitions can be easily obtained by modifying the analogous easy proof for Hopf algebras.

7. THE QUANTUM DOUBLE OF A SEMISIMPLE T-COALGEBRA

The quantum double of a semisimple Hopf algebra over a field of characteristic zero is both semisimple and modular (see [3]). We start this section by recalling the definition of a semisimple T-coalgebra [22], a modular Hopf algebra [14], and a modular T-coalgebra [20, 22]. After that, given any totally-finite T-coalgebra H , we discuss the relation between $D(H)$ and the quantum double of H_{pk} . Finally, we discuss the semisimplicity and the modularity of the quantum double $D(H)$ of a semisimple T-coalgebra H over a field of characteristic zero. In particular, we prove that $D(H)$ is semisimple if and only if H is totally-finite. Moreover, when H is totally-finite, $D(H)$ is also modular.

BASIC DEFINITIONS. Let H be a T-coalgebra. We say that H is *semisimple* when any algebra H_{α} (with $\alpha \in \pi$) is semisimple. It is proved in [22] that H is semisimple if and only if H_1 is semisimple. Further, by [18], infinite-dimensional Hopf algebras over a field are never semisimple. It follows that a necessary condition for H to be semisimple is that H_1 is finite-dimensional.

Let $H_1 = (H_1, R_1 = \xi_{1.i} \otimes \zeta_{1.i}, \theta_1)$ be a ribbon Hopf algebra. Given a finite-dimensional representation V of H_1 , and a H -linear endomorphism $f: V \rightarrow V$, the *quantum trace* $\text{trq}(f)$ of f is defined as $\text{trq}(f) = \text{tr}(u_1\theta_1 f)$, where $u_1 = s_1(\zeta_{1.i})\xi_{1.i}$ and $\text{tr}(\cdot)$ is the usual trace of endomorphisms. V is said to be *negligible* when $\text{trq}(\text{Id}_V) = 0$.

A *modular Hopf algebra* H_1 is a ribbon Hopf algebra endowed with a finite family of simple finite-dimensional H_1 -modules $\{V_i\}_{i \in I}$ satisfying the following conditions.

- There exists $0 \in I$ such that $V_0 = \mathbb{k}$ (with the structure of H_1 -module given by the comultiplication).
- For any $i \in I$, there exists $i^* \in I$ such that V_{i^*} is isomorphic to V_i^* .
- For any $j, k \in I$, the H_1 -module $V_j \otimes V_k$ is isomorphic to a finite sum of certain elements of $\{V_i\}_{i \in I}$, possibly with repetitions, and a negligible H_1 -module.

- Let $(S_{i,j})_{i,j \in I}$ denote the square matrix whose entry $S_{i,j}$ is the quantum trace of the endomorphism $x \mapsto \zeta_{1,k} \xi_{1,l} \otimes \xi_{1,k} \zeta_{1,l} x$ of $V_i \otimes V_j^*$. Then $S[H]$ is invertible.

A *modular T-coalgebra* [20] is a ribbon T-coalgebra H such that its component H_1 is a modular Hopf algebra.

Theorem 7.1. *The quantum double $D(H)$ of a T-coalgebra H over a field of characteristic 0 is semisimple if and only if H is totally-finite, and, in that case, $D(H)$ is also modular.*

To prove Theorem 7.1 we firstly need to discuss how the quantum double of a totally-finite T-coalgebra H and the quantum double of the packed form of H are related.

THE STRUCTURE OF A TOTALLY FINITE T-COALGEBRA. When H is a totally-finite T-coalgebra, the Hopf algebra $H_1^{\text{tot}} = (H^*)_{\text{pk}}$ is the dual of a certain Hopf algebra H_{pk} . An easy computation shows that H_{pk} satisfies the following conditions.

- As an algebra, $H_{\text{pk}} = \bigoplus_{\alpha \in \pi} H_\alpha$.
- The comultiplication Δ_{pk} is obtained setting $\Delta_{\text{pk}} = \sum_{\beta, \gamma: \beta\gamma=\alpha} \Delta_{\beta, \gamma}(h)$, for any $h \in H_\alpha \subset H_{\text{pk}}$.
- The counit ε_{pk} is given by $\varepsilon_{\text{pk}}|_{H_1} = \varepsilon$ and $\varepsilon_{\text{pk}}|_{H_\alpha} = 0$, for any $\alpha \in \pi$, $\alpha \neq 1$.
- The antipode is given by $s_{\text{pk}} = \sum_{\alpha \in \pi} s_\alpha$.

Moreover, we have a group homomorphism $\varphi_{\text{pk}}: \pi \rightarrow \text{Aut}(H_{\text{pk}}): \alpha \mapsto \bigoplus_{\beta \in \pi} \varphi_\beta^\alpha$.

Conversely, let H_{tot} be a finite-dimensional Hopf algebra (with antipode s_{tot} , counit ε and comultiplication Δ_{tot}), endowed with a family of subcoalgebras $\{H_\alpha\}_{\alpha \in \pi}$ and a group homomorphism $\varphi_{\text{tot}}: \pi \rightarrow \text{Aut}(H_{\text{tot}}): \alpha \mapsto \varphi_{\text{tot}, \alpha}$, such that the following conditions hold.

- H_{tot} is, as an algebra, the product H_α ($\alpha \in \pi$).
- For any $\alpha \in \pi$, we have $\Delta_{\text{tot}}(H_\alpha) \subset \bigoplus_{\beta, \gamma \text{ s.t. } \beta\gamma=\alpha} (H_\beta \otimes H_\gamma)$.
- For any $\alpha \in \pi \setminus \{1\}$, $H_\alpha \subset \text{Ker } \varepsilon$.
- For any $\alpha \in \pi$, $s_{\text{tot}}(H_\alpha) = H_{\alpha^{-1}}$.
- For any $\alpha, \beta \in \pi$, the image of H_α under $\varphi_{\text{tot}, \beta}$ lies in $H_{\beta\alpha\beta^{-1}}$.

To H_{tot} corresponds, in the obvious way, a T-coalgebra H such that $H_{\text{pk}} = H_{\text{tot}}$. In particular, for any $\alpha, \beta \in \pi$, the component $\Delta_{\alpha, \beta}: H_{\alpha\beta} \rightarrow H_\alpha \otimes H_\beta$ of the comultiplication is given by $H_{\alpha\beta} \hookrightarrow H_{\text{tot}} \xrightarrow{\Delta_{\text{tot}}} H_{\text{tot}} \otimes H_{\text{tot}} \xrightarrow{p_\alpha \otimes p_\beta} H_\alpha \otimes H_\beta$, where p_α and p_β are the canonical projections of H_{tot} on H_α and, respectively, H_β .

THE QUANTUM DOUBLE OF A TOTALLY-FINITE T-COALGEBRA. Let H be a totally-finite T-coalgebra. We have seen that $D(H)$ is totally finite. We can construct in the usual way the quantum double $D(H_{\text{pk}})$ of H_{pk} . Neither as an algebra nor as a coalgebra $D(H_{\text{pk}})$ is isomorphic to $(D(H))_{\text{pk}}$ since neither the multiplication nor the comultiplication of $D(H_{\text{pk}})$ depend on the conjugation.

Now, the canonical embedding of vector spaces $H_1 \otimes H_1^{\text{tot}, \text{cop}} \hookrightarrow \bigoplus_{\alpha \in \pi} H_\alpha \otimes H_1^{\text{tot}, \text{cop}}$ provides an embedding of Hopf algebras of $D_1(H) \hookrightarrow D(H_{\text{pk}})$, so we can identify $D_1(H)$ with its image in $D(H_{\text{pk}})$. Moreover, even if the universal R -matrix $R_{1,1} = (e_{1,i} \otimes \varepsilon) \otimes (1 \otimes e^{1,i})$ of $D_1(H)$ and the universal R -matrix $R_{\text{pk}} = \sum_{\alpha \in \pi} R_{\alpha, \alpha}$ of $D(H_{\text{pk}})$ are different, for any $x \in H_1 \otimes H^*$ we have

$$(32) \quad xR_{\text{pk}} = xR_{1,1} \quad \text{and} \quad R_{\text{pk}}x = R_{1,1}x.$$

FACTORIZABLE T-COALGEBRAS. We recall that a finite dimensional quasitriangular Hopf algebra H with universal R -matrix $R = \xi_i \otimes \zeta_i$ is *factorizable* (see [12]) if the map $\lambda: H^* \rightarrow H: f \mapsto \langle f, \zeta_i \xi_j \rangle \xi_i \zeta_j$ is bijective. In particular, the quantum double of any finite-dimensional Hopf algebra is factorizable. It is proved in [11] that any factorizable Hopf algebra is unimodular.

Lemma 7.2. *Let H be a finite-type T-coalgebra. The Hopf algebra $(D_1(H), \xi_{(1).i} \otimes \zeta_{(1).i})$ is factorizable.*

Proof. We can identify $D_1(H)$ as a subspace of $D(H_{\text{pk}})$ and $(D_1(H))^*$ as a subspace of $(D(H_{\text{pk}}))^*$. We only need to check that $\lambda_{D_1(H)} = \lambda_{D(H_{\text{pk}})}|_{(D_1(H))^*}$. Let $R = \xi_i \otimes \zeta_i$ be the universal R -matrix of $D(H_{\text{pk}})$. For any $f \in (D_1(H))^*$ we have

$$\lambda_{D(H_{\text{pk}})}(f) = \langle f, \zeta_i \xi_j \rangle \xi_i \zeta_j = \langle f, \zeta_{(1).i} \xi_{(1).j} \rangle \xi_{(1).i} \zeta_{(1).j} = \lambda_{D_1(H)}(f).$$

□

THE QUANTUM DOUBLE OF A SEMISIMPLE T-COALGEBRA. Let us consider the case of a semisimple T-coalgebra H over a field $\mathbb{k}$ of characteristic zero. It was proved in [22] that, for any $\alpha \in \pi$,

$$(33) \quad s_{\alpha^{-1}} \circ s_{\alpha} = \text{Id}_{H_{\alpha}}.$$

Lemma 7.3. *If H is quasitriangular, then it is also ribbon by setting $\theta_{\alpha} = u_{\alpha}^{-1}$, for any $\alpha \in \pi$.*

Proof. Axiom 31a follows by (27g) and (33). Axiom 31c follows by (30b). Axiom 31d follows by (27f). Axiom 31e can be rewritten

$$(34) \quad s_{\alpha^{-1}}(u_{\alpha^{-1}}) = u_{\alpha^{-1}}.$$

This follows by [22, Theorem 6(b)], by observing that, in that formula, $g_{\alpha} = 1_{\alpha}$ (by [22, Corollary 7]) $\hat{\varphi}(\alpha) = 1$ (by [22, Theorem 7]), and $h_{\alpha} = 1_{\alpha}$ (by [22, Lemma 16], since the distinguished group-like element of H_1^* is equal to ε because H_1^* is semisimple by [7]). Finally axiom 31b follows by (34). □

Proof of Theorem 7.1. If H is not totally-finite, $D_1(H)$ is not finite-dimensional and so it is not semisimple.

Suppose that H is totally-finite. H_{α} is semisimple for any $\alpha \in \pi$, therefore H_{pk} is semisimple. It follows that $D(H_{\text{pk}})$ is semisimple (see [11]). Since $D_1(H)$ can be identified with a subalgebra of $D(H_{\text{pk}})$, also $D_1(H)$ is semisimple, hence $D(H)$ is semisimple. By Lemma 7.3, $D(H)$ has a natural structure of a ribbon T-coalgebra.

The rest of the proof follows the streamline of the proof of Lemma 1.1 in [3] to which we refer for further details. Let $C(D_1(H)) \subset (D_1(H))^*$ be the ring of characters of $D_1(H)$ and let $Z(D_1(H))$ be the center of $D_1(H)$. By Lemma 7.2, the restriction $\Lambda = \lambda|_{C(D_1(H))} : C(D_1(H)) \rightarrow Z(D_1(H))$ is an isomorphism of algebras. Let $\text{Irr}(D_1(H)) = \{V_i | 0 \leq i \leq m\}$ be a set of representatives for the isomorphism classes of the irreducible representations of $D(H_1)$ such that $V_0 = \mathbb{k}$. Then $B = \{\chi_{j^*} = \text{tr}|_{V_{j^*}} : 0 \leq j \leq m\}$ is a linear basis of $C(D(H_1))$. We also observe that the set $C = \{e_j : 0 \leq j \leq m\}$ of central primitive idempotents of $D_1(H)$ is a basis of $Z(D(H_1))$. Now, $S[D_1(H)] = DA$, where A is the invertible matrix which represents Λ with respect to the bases B and C , while $D = \text{diag}(\dim V_i)$. □

8. THE RIBBON EXTENSION OF A QUASITRIANGULAR T-COALGEBRA

Let H be any quasitriangular T-coalgebra (not necessarily of finite-type). We describe how to obtain a ribbon T-coalgebra $\text{RT}(H)$ such that, when $\pi = \{1\}$, we recover the construction described in [13].

DEFINITION OF $RT(H)$. The *ribbon extension* $RT(H)$ of a quasitriangular T-coalgebra H is the T-coalgebra defined as follows.

- For any $\alpha \in \pi$, the α -th component of $RT(H)$, denoted $RT_\alpha(H)$, is the vector space whose elements are formal expressions $h + kv_\alpha$, with $h, k \in H_\alpha$. The sum is given by $(h + kv_\alpha) + (h' + k'v_\alpha) = (h + h') + (k + k')v_\alpha$, for any $h, h', k, k' \in H_\alpha$. The multiplication is obtained by requiring $v_\alpha^2 = u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}})$, i.e., by setting, for any $h, h', k, k' \in H_\alpha$,

$$(h + kv_\alpha)(h' + k'v_\alpha) = (hh' + k\varphi_\alpha(k')u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}})) + (hk' + k\varphi_\alpha(k'))v_\alpha.$$

We identify H_α with the subset $\{h + 0v_\alpha | h \in H_\alpha\}$ of $RT_\alpha(H)$. The algebra $RT_\alpha(H)$ is unitary with unit $1_\alpha = 1_\alpha + 0v_\alpha$. Moreover, for any $\alpha, \beta \in \pi$, we have $R_{\alpha,\beta} \in H_\alpha \otimes H_\beta \subset RT_\alpha(H) \otimes RT_\beta(H)$.

- The comultiplication is given by

$$\Delta_{\alpha,\beta}(h + kv_\alpha) = \Delta_{\alpha,\beta}(h) + \Delta_{\alpha,\beta}(k)\tilde{Q}_{\alpha,\beta}(v_\alpha \otimes v_\beta),$$

for any $h, k \in H_\alpha$ and $\alpha, \beta \in \pi$. The counit is given by $\langle \varepsilon, h + kv_\alpha \rangle = \langle \varepsilon, h \rangle + \langle \varepsilon, k \rangle$, for any $h, k \in H_1$.

- The antipode is given by $s_\alpha(h + kv_\alpha) = s_\alpha(h) + (s_\alpha \circ \varphi_{\alpha^{-1}})(k)v_{\alpha^{-1}}$, for any $h, k \in H_\alpha$ and $\alpha \in \pi$.
- Finally, the conjugation is given by $\varphi_\beta(h + kv_\alpha) = \varphi_\beta(h) + \varphi_\beta(k)v_{\beta\alpha\beta^{-1}}$, for any $h, k \in H_\alpha$ and $\alpha, \beta \in \pi$.

Theorem 8.1. $RT(H)$ is a ribbon T-coalgebra.

Proof. Since computations involved to prove in detail Theorem 8.1 are relatively long, we only prove the coassociativity of Δ , while the rest is left to the reader. To prove the multiplicativity of Δ you need to check before the relation

$$\Delta_{\alpha,\beta}(u_{\alpha\beta}s_{(\alpha\beta)^{-1}}(u_{(\alpha\beta)^{-1}})) = \tilde{Q}_{\alpha,\beta}(\varphi_\alpha \otimes \varphi_\beta)(\tilde{Q}_{\alpha,\beta}u_\alpha s_{\alpha^{-1}}(u_{\alpha^{-1}}) \otimes u_\beta s_{\beta^{-1}}(u_{\beta^{-1}})).$$

(for any $\alpha, \beta \in \pi$). To prove that $s_{RT(H)}$ is an antipode, one first needs to show that it is antimultiplicative.

COASSOCIATIVITY. We need to check that, for any $h, k \in H_{\alpha\beta\gamma}$, with $\alpha, \beta, \gamma \in \pi$, we have

$$(35) \quad ((\Delta_{\alpha,\beta} \otimes RT_\gamma(H)) \circ \Delta_{\alpha\beta,\gamma})(h + kv_{\alpha\beta\gamma}) = ((RT_\alpha(H) \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma})(h + kv_{\alpha\beta\gamma}).$$

By the linearity and the multiplicativity of the comultiplication in $RT(H)$ and the coassociativity of the comultiplication in H , by computing the left-hand side of (35), we obtain $((\Delta_{\alpha,\beta} \otimes RT_\gamma(H)) \circ \Delta_{\alpha\beta,\gamma})(h + kv_{\alpha\beta\gamma}) = h'_{(\alpha)} \otimes h''_{(\beta)} \otimes h'''_{(\gamma)} + (k'_{(\alpha)} \otimes k''_{(\beta)} \otimes k'''_{(\gamma)})((\Delta_{\alpha,\beta} \otimes RT_\gamma(H)) \circ \Delta_{\alpha\beta,\gamma})(v_{\alpha\beta\gamma})$, while, by computing the right-hand side we obtain $((RT_\alpha(H) \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma})(h + kv_{\alpha\beta\gamma}) = h'_{(\alpha)} \otimes h''_{(\beta)} \otimes h'''_{(\gamma)} + (k'_{(\alpha)} \otimes k''_{(\beta)} \otimes k'''_{(\gamma)})((RT_\alpha(H) \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma})(v_{\alpha\beta\gamma})$, so we only need to check that we have

$$(36) \quad ((\Delta_{\alpha,\beta} \otimes RT_\gamma(H)) \circ \Delta_{\alpha\beta,\gamma})(v_{\alpha\beta\gamma}) = ((RT_\alpha(H) \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma})(v_{\alpha\beta\gamma}).$$

By computing the left-hand side of (36), we get $((\Delta_{\alpha,\beta} \otimes RT_\gamma(H)) \circ \Delta_{\alpha\beta,\gamma})(v_{\alpha\beta\gamma}) = ((\Delta_{\alpha,\beta} \otimes H_\gamma)(\tilde{Q}_{\alpha\beta,\gamma}))(\tilde{Q}_{\alpha,\beta})_{12\gamma}(v_\alpha \otimes v_\beta \otimes v_\gamma)$, while, by computing the right-hand side, $((RT_\alpha(H) \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma})(v_{\alpha\beta\gamma}) = ((H_\alpha \otimes \Delta_{\beta,\gamma})(\tilde{Q}_{\alpha,\beta\gamma}))(\tilde{Q}_{\beta,\gamma})_{\alpha 23}(v_\alpha \otimes v_\beta \otimes v_\gamma)$. We only need to show that $((\Delta_{\alpha,\beta} \otimes H_\gamma)(\tilde{Q}_{\alpha\beta,\gamma}))(\tilde{Q}_{\alpha,\beta})_{12\gamma} = ((H_\alpha \otimes \Delta_{\beta,\gamma})(\tilde{Q}_{\alpha,\beta\gamma}))(\tilde{Q}_{\beta,\gamma})_{\alpha 23}$.

$\Delta_{\beta,\gamma})(\tilde{Q}_{\alpha,\beta\gamma}))(\tilde{Q}_{\beta,\gamma})_{\alpha 23}$, or, equivalently,

$$\begin{aligned}
 (37) \quad & \left((\sigma \circ (H_\beta \otimes \varphi_\alpha))(R_{\beta,\alpha}) \right)_{12\gamma} (R_{\alpha,\beta})_{12\gamma} \cdot \\
 & \cdot ((\Delta_{\alpha,\beta} \otimes H_\gamma) \circ \sigma \circ (H_\gamma \otimes \varphi_{\alpha\beta}))(R_{\gamma,\alpha\beta})(\Delta_{\alpha,\beta} \otimes H_\gamma)(R_{\alpha\beta,\gamma}) \\
 & = \left((\sigma \circ (H_\gamma \otimes \varphi_\beta))(R_{\gamma,\beta}) \right)_{\alpha 23} (R_{\beta,\gamma})_{\alpha 23} \cdot \\
 & \cdot ((H_\alpha \otimes \Delta_{\beta,\gamma}) \circ \sigma \circ (H_{\beta\gamma} \otimes \varphi_\alpha))(R_{\beta\gamma,\alpha})(H_\alpha \otimes \Delta_{\beta,\gamma})(R_{\alpha,\beta\gamma}).
 \end{aligned}$$

Let us set $x = \left((\sigma \circ (\varphi_{\alpha^{-1}} \otimes H_\alpha) \circ \Delta_{\alpha\beta\alpha^{-1},\alpha}) \otimes H_\gamma \right) \circ \sigma \circ (H_\gamma \otimes \varphi_{\alpha\beta}) (R_{\gamma,\alpha\beta})$, and $y = \left((H_\alpha \otimes (\sigma \circ (\varphi_{\beta^{-1}} \otimes H_\beta) \circ \Delta_{\beta\gamma\beta^{-1},\beta})) \circ \sigma \circ (H_{\beta\gamma} \otimes \varphi_\alpha) \right) (R_{\beta\gamma,\alpha})$. Since, by (7a), $(R_{\alpha,\beta})_{12\gamma} ((\Delta_{\alpha,\beta} \otimes H_\gamma) \circ \sigma \circ (H_\gamma \otimes \varphi_{\alpha\beta}))(R_{\gamma,\alpha\beta}) = x(R_{\alpha,\beta})_{12\gamma}$ and $(R_{\beta,\gamma})_{\alpha 23} ((H_\alpha \otimes \Delta_{\beta,\gamma}) \circ \sigma \circ (H_{\beta\gamma} \otimes \varphi_\alpha))(R_{\beta\gamma,\alpha}) = y(R_{\beta,\gamma})_{\alpha 23}$, if we substitute these expressions in (37) and we apply axiom (7c) to the left-hand side and axiom (7b) to the right-hand side, we find that (37) can be rewritten as

$$\begin{aligned}
 & \left((\sigma \circ (H_\beta \otimes \varphi_\alpha))(R_{\beta,\alpha}) \right)_{12\gamma} x(R_{\alpha,\beta})_{12\gamma} ((H_\alpha \otimes \varphi_{\beta^{-1}})(R_{\alpha,\beta\alpha\beta^{-1}}))_{1\beta 3} (R_{\beta,\gamma})_{\alpha 23} \\
 & = \left((\sigma \circ (H_\gamma \otimes \varphi_\beta))(R_{\gamma,\beta}) \right)_{\alpha 23} y(R_{\beta,\gamma})_{\alpha 23} (R_{\beta,\gamma})_{\alpha 23} (R_{\alpha,\gamma})_{1\beta 3} (R_{\alpha,\beta})_{12\gamma}.
 \end{aligned}$$

Thus, by the Yang-Baxter equation (8), we can rewrite (37) as

$$(38) \quad \left((\sigma \circ (H_\beta \otimes \varphi_\alpha))(R_{\beta,\alpha}) \right)_{12\gamma} x = \left((\sigma \circ (H_\gamma \otimes \varphi_\beta))(R_{\gamma,\beta}) \right)_{\alpha 23} y.$$

Given three vector spaces V_1 , V_2 , and V_3 , let us introduce the notation $\sigma_{i,j,k}$ (with $\{i,j,k\} = \{1,2,3\}$) for the permutation $V_1 \otimes V_2 \otimes V_3 \rightarrow V_i \otimes V_j \otimes V_k$.

If we compute the two factors on the left-hand side in (38), then we have

$$\begin{aligned}
 & \left((\sigma \circ (H_\beta \otimes \varphi_\alpha))(R_{\beta,\alpha}) \right)_{12\gamma} = ((\sigma \otimes H_\gamma) \circ \sigma_{2,3,1} \circ (H_\gamma \otimes H_\beta \otimes \varphi_\alpha))((R_{\beta,\alpha})_{\gamma 23}) \\
 & = ((\sigma \otimes H_\gamma) \circ \sigma_{2,3,1} \circ (H_\beta \otimes \varphi_\beta \otimes \varphi_{\alpha\beta}))((R_{\beta,\beta^{-1}\alpha\beta})_{\gamma 23}),
 \end{aligned}$$

(where in the last passage we used axiom (7d)) and

$$\begin{aligned}
 x & = \left(\left((\sigma \circ (\varphi_{\alpha^{-1}} \otimes H_\alpha) \circ \Delta_{\alpha^{-1}\beta\alpha,\alpha} \circ \varphi_{\alpha\beta}) \otimes H_\gamma \right) \circ \sigma \right) (R_{\gamma,\alpha\beta}) \\
 & = \left(\left((\sigma \circ (\varphi_{\alpha^{-1}} \otimes H_\alpha) \circ (\varphi_{\alpha\beta} \otimes \varphi_{\alpha\beta})) \otimes H_\gamma \right) \circ \sigma_{2,3,1} \circ (H_\gamma \otimes \Delta_{\beta,\beta^{-1}\alpha\beta}) \right) (R_{\gamma,\alpha\beta}) \\
 & = ((\sigma \otimes H_\gamma) \circ \sigma_{2,3,1} \circ (H_\beta \otimes \varphi_\beta \otimes \varphi_{\alpha\beta}))((R_{\gamma,\beta^{-1}\alpha\beta})_{1\beta 3} (R_{\gamma,\beta})_{12(\beta^{-1}\alpha\beta)}),
 \end{aligned}$$

(where, in the last passage, we used axiom (7b)). Similarly, on the right-hand side we have

$$\left((\sigma \circ (H_\gamma \otimes \varphi_\beta))(R_{\gamma,\beta}) \right)_{\alpha 23} = ((H_\alpha \otimes \sigma) \circ \sigma_{3,1,2} \circ (H_\gamma \otimes \varphi_\beta \otimes \varphi_{\alpha\beta}))((R_{\gamma,\beta})_{12(\beta^{-1}\alpha\beta)})$$

and

$$\begin{aligned}
y &= \left((H_\alpha \otimes \sigma) \circ \left(\varphi_\alpha \otimes ((\varphi_{\beta^{-1}} \otimes H_\beta) \circ \Delta_{\beta\gamma\beta^{-1},\beta}) \right) \circ \sigma \right) (R_{\beta\gamma,\alpha}) \\
&= \left((H_\alpha \otimes \sigma) \circ (\varphi_{\alpha\beta} \otimes H_\gamma \otimes \varphi_\beta) \circ (\varphi_{\beta^{-1}} \otimes (\Delta_{\gamma,\beta} \circ \varphi_{\beta^{-1}})) \circ \sigma \right) (R_{\beta\gamma,\alpha}) \\
&\text{(by axiom (7d))} \\
&= ((H_\alpha \otimes \sigma) \circ (\varphi_{\alpha\beta} \otimes H_\gamma \otimes \varphi_\beta) \circ (H_{\beta^{-1}\alpha\beta} \otimes \Delta_{\gamma,\beta}) \circ \sigma) (R_{\gamma\beta,\beta\alpha\beta^{-1}}) \\
&= ((H_\alpha \otimes \sigma) \circ (\varphi_{\alpha\beta} \otimes H_\gamma \otimes \varphi_\beta) \circ \sigma_{3,1,2} \circ (\Delta_{\gamma,\beta} \otimes H_{\beta^{-1}\alpha\beta})) (R_{\gamma\beta,\beta\alpha\beta^{-1}}) \\
&= ((H_\alpha \otimes \sigma) \circ \sigma_{3,1,2} \circ (H_\gamma \otimes \varphi_\beta \otimes \varphi_{\alpha\beta})) \left(((H_\gamma \otimes \varphi_{\beta^{-1}})(R_{\gamma,\alpha}))_{1\beta 3} (R_{\beta,\beta^{-1}\alpha\beta})_{\gamma 23} \right)
\end{aligned}$$

(where in the last passage we used axiom (7c)).

We observe that the application $(H_\alpha \otimes \sigma) \circ \sigma_{3,1,2} \circ (H_\alpha \otimes \varphi_\alpha \otimes \varphi_{\alpha\beta})$ is bijective. So, we can rewrite (38) in the form

$$\begin{aligned}
&(R_{\beta,\beta^{-1}\alpha\beta})_{\gamma 23} (R_{\gamma,\beta^{-1}\alpha\beta})_{1\beta 3} (R_{\gamma,\beta})_{12(\beta^{-1}\alpha\beta)} \\
&= (R_{\gamma,\beta})_{12(\beta^{-1}\alpha\beta)} ((H_\gamma \otimes \varphi_{\beta^{-1}})(R_{\gamma,\alpha}))_{1\beta 3} (R_{\beta,\beta^{-1}\alpha\beta})_{\gamma 23}
\end{aligned}$$

and this last formula is true by the Yang-Baxter equation (8). $\square$

Corollary 8.2. *Let H be a finite-type T-coalgebra. We obtain a ribbon T-coalgebra $\text{RT}(D(H))$.*

REFERENCES

1. Eiichi Abe, *Hopf Algebras*, Cambridge Tractats in Mathematics 74, Cambridge University Press, Cambridge (U.K.), 1980. MR **83a**:16010
2. Vladimir G. Drinfeld, *Quantum groups*, Proceedings of the International Congress of Mathematicians (Berkeley, 1986) (Providence, RI), vol. 1, Amer. Math. Soc., 1987, pp. 798–820. MR **89f**:17017
3. Pavel Etingof and Shlomo Gelaki, *Some properties of finite-dimensional semisimple Hopf algebras*, Math. Res. Lett. **5** (1998), 191–197. MR **99e**:16050
4. André Joyal and Ross Street, *Braided tensor categories*, Adv. in Math. **102** (1993), no. 1, 20–78. MR **94m**:18008
5. Christian Kassel, *Quantum Groups*, Graduate Texts in Mathematics 155, Springer-Verlag, New York, 1995. MR **96e**:17041
6. Christian Kassel and Vladimir G. Turaev, *Double construction for monoidal categories*, Acta Math. **175** (1995), no. 1, 1–48. MR **96m**:18015
7. Richard G. Larson and David E. Radford, *Semisimple cosemisimple Hopf algebras*, Amer. J. Math. **110** (1988), no. 1, 187–195. MR **89a**:16011
8. Thang Le and Vladimir G. Turaev, *Quantum groups and ribbon G-categories*, Preprint math.QA/0103017, 2001.
9. Shahn Majid, *Physics for algebraists: noncommutative and noncocommutative Hopf algebras by a bicrossproduct construction*, J. Algebra **130** (1990), no. 1, 17–64. MR **91j**:16050
10. ———, *Representations, duals and quantum doubles of monoidal categories*, Proceedings of the Winter School on Geometry and Physics (Srni, 1990), no. 26, 1991, pp. 197–206. MR **93c**:18008
11. David E. Radford, *Minimal quasitriangular Hopf algebras*, J. Algebra **157** (1993), no. 2, 285–315. MR **94c**:16052
12. Nicolai Yu. Reshetikhin and Michael A. Semenov-Tian-Shansky, *Quantum R-matrices and factorization problems*, J. Geom. Phys. **5** (1988), no. 4, 533–550 (1989). MR **92g**:17019
13. Nicolai Yu. Reshetikhin and Vladimir G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, Comm. Math. Phys. **127** (1990), no. 1, 1–26. MR **91c**:57016
14. ———, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), no. 3, 547–597. MR **92b**:57024
15. Hans-Jürgen Schneider, *Lectures on Hopf Algebras*, Available at the site <http://www.mathematik.uni-muenchen.de/~hanssch/Publications.html>, 1995.

16. Ross Street, *The quantum double and related constructions*, J. Pure Appl. Algebra **132** (1998), no. 2, 195–206. MR **99e**:16054
17. Moss E. Sweedler, *Hopf Algebras*, Mathematics Lecture Note Series, W. A. Benjamin, Inc., New York, 1969. MR 40/5705
18. ———, *Integrals for Hopf algebras*, Ann. of Math. **89** (1969), no. 2, 323–335. MR 39/4167
19. Vladimir G. Turaev, *Homotopy field theory in dimension 2 and group-algebras*, Preprint **math.QA/9910010**, 2000.
20. ———, *Homotopy field theory in dimension 3 and crossed group-categories*, Preprint **math.GT/000529**, 2000.
21. Leonid Vainermann, In preparation.
22. Alexis Virelizier, *Hopf group-coalgebras*, Preprint **math.QA/0012073**. To appear in *J. Pure Appl. Algebra*, 2000.
23. ———, *Algèbres de Hopf graduées et fibrés plats sur le 3-variétés* (In English), Ph.D. thesis, Université Louis Pasteur (Strasbourg), 2001.
24. David N. Yetter, *Quantum groups and representations of monoidal categories*, Math. Proc. Cambridge Philos. Soc. **108** (1990), no. 2, 261–290. MR **91k**:16028
25. Marco Zunino, *Bicrossproduct of crossed group coalgebras*, In preparation.
26. ———, *Tannaka Theory for Turaev crossed structures*, In preparation.
27. ———, *Yang-Baxter modules for Turaev crossed structures*, In preparation.

MARCO ZUNINO – INSTITUT DE RECHERCHE MATHÉMATIQUES AVANCÉE (IRMA) – UNIVERSITÉ LOUIS PASTEUR ET CNRS – 7, RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX, FRANCE.

E-mail address: **zunino@math.u-strasbg.fr**